

Product Datasheet

PSTAIR Antibody (PSTAIR)

NB120-10345

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB120-10345

Updated 10/23/2016 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB120-10345




NB120-10345**PSTAIR Antibody (PSTAIR)**

Product Information	
Unit Size	0.1 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSTAIR
Preservative	0.1% Sodium Azide
Isotype	IgG1
Purity	Unpurified
Buffer	Raw ascites
Product Description	
Host	Mouse
Species	Human, Mouse, Avian, Plant, Primate, Xenopus
Reactivity Notes	Cross-reacts with human, monkey, mouse, Japanese quail, fish (goldfish, carp, eel, starfish, and amago salmon), Xenopus, ciliates, and plants (lily and onion). Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.
Specificity/Sensitivity	PSTAIR (PSTAIR)
Immunogen	Synthetic peptide: EGV PSTAIREISLLKE conjugated to BSA.
Product Application Details	
Applications	Western Blot, ELISA, Immunoprecipitation
Recommended Dilutions	Western Blot 1:100-1:2000, ELISA 1:100-1:2000, Immunoprecipitation 1:10-1:500
Application Notes	ELISA: Use at an assay dependent dilution. IP: Use at an assay dependent dilution. It cannot precipitate p34cdc2; when it is complexed with cyclin B.WB: Use at a dilution of 1/4000 (determined by indirect blotting using COS-7 cell extract). The antibody recognises 31-34kD proteins (1-4) bands. Monoclonal Anti-PSTAIR may be used for the localization of cyclin-dependent kinases, containing the PSTAIR motif. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

Images

Western Blot: PSTAIR Antibody (PSTAIR) [NB120-10345] - Cell line lysates were separated on SDS-PAGE and probed with 1:500 Monoclonal Anti-PSTAIR Clone:PSTAIR. The antibody was developed using Goat Anti-Mouse IgG-Peroxidase and a chemiluminescent substrate. Lanes 1. HEK-293T 2. HeLa 3. SK-N-SH 4. SH-SY5Y 5. U87 6. COS7 7. PC12 8. Neuro-2a 9. Arabidopsis





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Novus Biologicals Canada

461 North Service Road West, Unit B37
Oakville, ON L6M 2V5
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada@novusbio.com

Novus Biologicals Europe

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Products Related to NB120-10345

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005	Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB120-10345

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

