

Product Datasheet

MMP-2 Antibody (2C1) - (Pro and Active) - BSA Free NB200-113SS

Unit Size: 0.025 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB200-113

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB200-113



NB200-113SS

MMP-2 Antibody (2C1) - (Pro and Active) - BSA Free


Product Information	
Unit Size	0.025 ml
Concentration	1.0 mg/ml
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	2C1
Preservative	0.02% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Mouse MMP-2 Antibody (2C1) - (Pro and Active) - BSA Free (NB200-113) is a monoclonal antibody validated for use in IHC, WB, ELISA, Flow, ICC/IF and IP. Anti-MMP-2 Antibody: Cited in 6 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	4313
Gene Symbol	MMP2
Species	Human, Mouse, Rat, Guinea Pig
Specificity/Sensitivity	This is specific for pro and active MMP-2.
Immunogen	Activated recombinant human MMP2. [UniProt# P08253]

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation
Recommended Dilutions	Western Blot 2-5 ug/ml, Flow Cytometry, ELISA 1 ug/ml, Immunohistochemistry 5 ug/ml, Immunocytochemistry/ Immunofluorescence reported in scientific literature (PMID 30681163), Immunoprecipitation 1:10-1:500, Immunohistochemistry-Paraffin reported in scientific literature (PMID 24828425), Immunohistochemistry-Frozen 5 ug/ml

Images


Western Blot: MMP-2 Antibody (2C1) - (Pro and Active) [NB200-113] - Recombinant human MMP-2 expressed from CHO cells was separated on a 7.5 % gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 2.0 ug/ml anti-MMP-2 in blocking buffer and detected with an anti-mouse HRP secondary antibody using chemiluminescence.




Western Blot: MMP-2 Antibody (2C1) - (Pro and Active) [NB200-113] -
Samples: 1) Human Recombinant MMP-2, and 2) Schwann Cell Culture
Medium Run on 10% SDS-PAGE, followed by electroblot to
nitrocellulose and slot blot immunolabeling. Antibodies: primary - NB200-
113, secondary - anti-mouse-biotin. Detection: avidin-HRP. Chromagen:
DAB/H2O2.




Immunohistochemistry-Paraffin: MMP-2 Antibody (2C1) - (Pro and Active) [NB200-113] -
Analysis of a FFPE human breast cancer section using 5 ug/ml conc. of MMP2 antibody (clone 2C1) on a Bond Rx
autostainer (Leica Biosystems). The assay involved 20 minutes of heat
induced antigen retrieval (HIER) with 10mM sodium citrate buffer (pH
6.0) and endogenous peroxidase quenching using peroxide block. The
sections were incubated with primary antibody for 30 minutes. Bond
Polymer Refine Detection (Leica Biosystems) and DAB were used for
signal detection which followed counterstaining with hematoxylin. Whole
slide scanning and capturing of representative images (20X) were
performed using Aperio AT2 (Leica Biosystems). MMP2
immunoreactivity was observed in the cytoplasm of the cancer cells and
the stromal cells, and in the inter-cellular spaces (secreted MMP2). The
cancer cells present in the core of tumor areas were found to be negative
for MMP2 staining.




Immunohistochemistry-Paraffin: MMP-2 Antibody (2C1) - (Pro and Active) [NB200-113] -
Analysis of a FFPE human breast cancer section using 5 ug/ml conc. of MMP2 antibody (clone 2C1) on a Bond Rx
autostainer (Leica Biosystems). The assay involved 20 minutes of heat
induced antigen retrieval (HIER) with 10mM sodium citrate buffer (pH
6.0) and endogenous peroxidase quenching using peroxide block. The
sections were incubated with primary antibody for 30 minutes. Bond
Polymer Refine Detection (Leica Biosystems) and DAB were used for
signal detection which followed counterstaining with hematoxylin. Whole
slide scanning and capturing of representative images (20X) were
performed using Aperio AT2 (Leica Biosystems). This MMP2 antibody
generated an expected immunoreactivity of MMP2 protein in the
cytoplasm of most of the cancer cells and the stromal cells, and in the
inter-cellular spaces (secreted pool of MMP2). Staining was performed
by Histowiz.



Immunohistochemistry-Paraffin: MMP-2 Antibody (2C1) - (Pro and Active) [NB200-113] -
Upper image: MMP-2 FITC (green), DAPI (blue)
Lower image: phase contrast of human choroid plexus (Alzheimer's
disease). Image from verified customer review.



Flow Cytometry: MMP-2 Antibody (2C1) - (Pro and Active) [NB200-113] - Flow Cytometry: MMP-2 Antibody (2C1) [Alexa Fluor® 405] [NB200-113AF405] - CD4+ T cells were activated for 24h before being fixed and permeabilized to detect MMP-2. Image using the Alexa Fluor 405 form of this antibody. Image from verified customer review.



Publications

Xiang A, Peng Z, He H et al. The potential of brimonidine for myopia treatment: Targeting MMP-2 to regulate choroidal thickness and control eye growth Heliyon 2024-09-04 [PMID: 39309849]

Najjar R Raspberry Polyphenols Target Molecular Pathways of Heart Failure Thesis 2023-01-01

Yu T, Xie X, Wei H et al. Electroacupuncture Improves Choroidal Blood Flow to Inhibit the Development of Lens-Induced Myopia in Guinea Pigs Evidence-Based Complementary and Alternative Medicine 2022-06-24 [PMID: 35783515] (WB, Guinea Pig)

Schilling BK, Schusterman MA 2nd, Kim DY et al. Adipose-derived stem cells delay muscle atrophy after peripheral nerve injury in the rodent model Muscle Nerve 2019-01-26 [PMID: 30681163] (ICC/IF, Rat)

Scheede-Bergdahl C, Bergdahl A, Schjerling P et al. Exercise-induced regulation of matrix metalloproteinases in the skeletal muscle of subjects with type 2 diabetes Diab Vasc Dis Res. 2014-06-04 [PMID: 24903024] (WB, Human)

Details:

MMP2 antibody used for WB (1:1000 dilution) on human skeletal muscle lysates from type 2 diabetes patients (T2DM / NIDDM) and control subjects (Full Text has detailed method on tissue lysis). Antibody detected the 72 kDa band of pro-MMP-2 and 62 kDa band of active-MMP-2 (Figure 3)

Xu H, Zhang Y, Sun H et al. Effects of Acupuncture at GV20 and ST36 on the Expression of Matrix Metalloproteinase 2, Aquaporin 4, and Aquaporin 9 in Rats Subjected to Cerebral Ischemia/Reperfusion Injury. PLoS ONE. 2014-05-15 [PMID: 24828425] (IHC-P, WB, Rat)

Rork TH, Hadzimichalis NM, Kappil MA, Merrill GF. Acetaminophen attenuates peroxyntirite-activated matrix metalloproteinase-2-mediated troponin I cleavage in the isolated guinea pig myocardium. J Mol Cell Cardiol 40(4):553-61. 2006-04-01 [PMID: 16530785] (WB, Guinea Pig)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-
techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB200-113

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

