

Product Datasheet

FXR/NR1H4 Antibody - BSA Free NB400-153SS

Unit Size: 0.025 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 3 Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB400-153

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB400-153



NB400-153SS

FXR/NR1H4 Antibody - BSA Free

Product Information	
Unit Size	0.025 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Glycine, 0.15M NaCl
Target Molecular Weight	66 kDa


Product Description	
Description	Novus Biologicals Rabbit FXR/NR1H4 Antibody - BSA Free (NB400-153) is a polyclonal antibody validated for use in WB and ICC/IF. Anti-FXR/NR1H4 Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	9971
Gene Symbol	NR1H4
Species	Human, Mouse
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 29491146).
Immunogen	A synthetic peptide made to the C-terminus of the human FXR protein. [UniProt# Q96RI1]

Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1 - 4 ug/mL, Immunocytochemistry/ Immunofluorescence 1:100
Application Notes	The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.




Images


Western Blot: FXR/NR1H4 Antibody [NB400-153] - Total protein from human A549, U2OS cell and mouse 3T3 cells was separated on a 7.5% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 2.0 ug/ml anti-FXR (NB400-153) in block buffer and detected with an anti-rabbit HRP secondary antibody using NovaLume chemiluminescence detection reagent (NPB2-61915).




Immunocytochemistry/Immunofluorescence: FXR/NR1H4 Antibody [NB400-153] - A549 cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.5% Triton-X100. The cells were incubated with anti-FXR/NR1H4 at 2 ug/ml overnight at 4C and detected with an anti-rabbit Dylight 488 (Green) at a 1:500 dilution. Alpha tubulin (DM1A) NB100-690 was used as a co-stain at a 1:1000 dilution and detected with an anti-mouse Dylight 550 (Red) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.



Western Blot: FXR Antibody [NB400-153] - Detection of FXR in HeLa nuclear extract using NB400-153. ECL 30 minute detection.



Immunocytochemistry/Immunofluorescence: FXR/NR1H4 Antibody [NB400-153] - NIH3T3 cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.5% Triton-X100. The cells were incubated with anti-FXR/NR1H4 at 2 ug/ml overnight at 4C and detected with an anti-rabbit Dylight 488 (Green) at a 1:500 dilution. Alpha tubulin (DM1A) NB100-690 was used as a co-stain at a 1:1000 dilution and detected with an anti-mouse Dylight 550 (Red) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.



Publications

Bai Y, Zhang J, Li J, Liao M et al. Silibinin, a commonly used therapeutic agent for non-alcohol fatty liver disease, functions through upregulating intestinal expression of fibroblast growth factor 15/19 Br J Pharmacol 2024-06-05 [PMID: 38839561]

Malivindi R, Santoro M, De Rose D et al. Activated-Farnesoid X Receptor (FXR) expressed in human sperm alters its fertilizing ability Reproduction 2018-06-19 [PMID: 29921626] (Human)

Gallo-Ebert C, Francisco J, Liu HY et al. Mice lacking ARV1 have reduced signs of metabolic syndrome and non-alcoholic fatty liver disease. J. Biol. Chem. 2018-02-28 [PMID: 29491146] (WB, Mouse)

Xing Y, Saner-Amigh K, Nakamura Y et al. The farnesoid X receptor regulates transcription of 3beta-hydroxysteroid dehydrogenase type 2 in human adrenal cells. Mol Cell Endocrinol 2009-02-27 [PMID: 19059462] (Human)



Procedures

Western Blot Protocol for FXR/NR1H4 Antibody (NB400-153)

FXR/NR1H4 Antibody:

Western Blot Protocol

1. Perform SDS-PAGE (4-12%) on samples to be analyzed, loading 20 ug of total transfected protein per lane.
 2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
 3. Stain the blot using ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
 4. Rinse the blot in TBS for approximately 5 minutes.
 5. Block the membrane using 5% non-fat dry milk in TBS + 0.5% BSA for 1 hour.
 6. Dilute the rabbit anti-FXR primary antibody (NB 400-153) in blocking buffer and incubate 2 hours at room temperature.
 7. Wash the membrane in water for 5 minutes and apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
 8. Wash the blot in TBS containing 0.05-0.1% Tween-20 for 10-20 minutes.
 9. Wash the blot in type I water for an additional 10-20 minutes (this step can be repeated as required to reduce background).
 10. Apply the detection reagent of choice in accordance with the manufacturers instructions (Amersham's ECL is the standard reagent used at Novus Biologicals).
- Note: Tween-20 can be added to the blocking buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.

Immunocytochemistry/ Immunofluorescence Protocol for FXR/NR1H4 Antibody (NB400-153)

FXR/NR1H4 Antibody:

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB400-153

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

