

Product Datasheet

HSP27 [p Ser86] Antibody NSB536

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NSB536

Updated 6/27/2018 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NSB536



NSB536**HSP27 [p Ser86] Antibody**


Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Purity	Immunogen affinity purified
Buffer	Dulbecco's PBS (pH 7.3), 1.0 mg/ml and 50% Glycerol

Product Description	
Host	Rabbit
Gene ID	3315
Gene Symbol	HSPB1
Species	Mouse
Reactivity Notes	Mouse HSP25. Endogenous human HSP27 phosphorylated at serine 82 (HeLa cells treated with TNF-a) was weakly detected by this antibody.
Specificity/Sensitivity	HSP25 Phosphospecific [Ser86]
Immunogen	The antiserum was produced against a chemically synthesized phosphopeptide derived from a region of mouse HSP25 that contains serine 86.

Product Application Details	
Applications	Western Blot
Recommended Dilutions	Western Blot
Application Notes	The antibody has been used for Western blotting 1:1000. The optimal antibody concentration should be determined empirically for each specific application.

Images

Peptide Competition Lysates prepared from NIH3T3 cells left unstimulated (1) or treated with anisomycin (2-6) were resolved by SDS-PAGE on a 10% polyacrylamide gel and transferred to PVDF. Membranes were left untreated (1-5) or treated with lambda (l) phosphatase (6), blocked with a 5% BSA-TBST buffer for one hour at room temperature, and incubated with HSP25 [pS86] (mouse) antibody for one hour at room temperature in 3% BSA-TBST buffer, following prior incubation with: no peptide (1, 2, 6), the non-phosphopeptide corresponding to the immunogen (3), a generic phosphoserine-containing peptide (4), or, the phosphopeptide immunogen (5). After washing, membranes were incubated with goat F(ab)2 anti-rabbit IgG HRP conjugate in 3% BSA-TBST buffer, and bands were detected using the Pierce SuperSignal(TM) method. The data show that only the peptide corresponding to HSP25 [pS86] (mouse) blocks the antibody signal, thereby demonstrating the specificity of the antibody. The signal was completely removed by l phosphatase treatment demonstrating that the antibody interacts specifically with the phosphorylated protein.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Novus Biologicals Canada

461 North Service Road West, Unit B37
Oakville, ON L6M 2V5
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada@novusbio.com

Novus Biologicals Europe

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NSB536

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

