

Rabbit Polyclonal anti-HIF-1 alpha, Sample Size

Catalog Number:	NB100-134SS
Background:	<p>Hypoxia contributes significantly to the pathophysiology of major categories of human disease, including myocardial and cerebral ischemia, cancer, pulmonary hypertension, congenital heart disease, and chronic obstructive pulmonary disease.</p> <p>HIF-1 is a nuclear protein involved in mammalian oxygen homeostasis. This occurs as a posttranslational modification by prolyl hydroxylation. HIF-1 is a heterodimer composed of HIF-1 alpha and HIF-1 beta subunits. Both subunits are constantly translated. However, under normoxic conditions, human HIF-1 alpha is hydroxylated at Pro402 or Pro564 by a set of HIF prolyl hydroxylases, is polyubiquitinated, and eventually degraded in proteosomes. Under hypoxic conditions, the lack of hydroxylation prevents HIF degradation and increases transcriptional activity. Therefore, the concentration of HIF-1 alpha increases in the cell. In contrast, HIF-1 beta remains stable under either condition. HIF hydroxylases provide insight into hypoxic cell responses, which may be used to help isolate therapeutic targets.</p>
Alternate Names:	anti-Hypoxia-inducible factor 1 alpha antibody, anti-HIF1 alpha antibody, anti-ARNT interacting protein antibody, anti-Hif1a antibody, anti-ARNT interacting protein antibody, anti-HIF-1alpha antibody, anti-Hypoxia inducible factor 1 alpha antibody, anti-Hypoxia inducible factor 1 alpha subunit basic helix antibody
Host:	Rabbit
Research Areas:	Hypoxia Inducible Factors and Prolyl-Hydroxylases, IHC Grade, Cancer Research
Immunogen:	Fusion protein containing amino acids 432-528 of human HIF-1 alpha.
Specificity:	This antibody is specific for HIF-1 alpha.
Localization:	Nucleus
Species Reactivity:	Human, mouse and rat. Mouse results mixed. Other species have not been tested.
Uses:	<p>This antibody is useful for ChIP, immunohistochemistry, and Western Blot. IHC applications have been mixed on mouse.</p> <p>* Other applications have not been tested.</p>
Dilutions:	<p>Suggested working dilutions *</p> <p>chromatin immunoprecipitation - use 25 ul antibody for 15 million cells (nuclear extracts recommended),</p> <p>immunohistochemistry 1:100,</p> <p>Western Blot 1:500-1:1,000,</p> <p>* Investigator should determine optimal working dilutions.</p>
Positive Controls:	<p>PC12 nuclear extracts (hypoxically induced), Cobalt Chloride treated Cos-7 nuclear extracts</p> <p>* Cos-7 Hypoxia Induced Nuclear Extract Kit NB800-PC26</p> <p>* PC12 Nuclear Extracts NB810-55229</p>
Packaging:	0.025 ml Whole Rabbit antisera.

- Preservative:** 0.05% Sodium azide
- Storage:** Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
- Notes:** You may use COS-7 treated and untreated nuclear extracts for your positive and negative controls for hypoxic upregulation (NB 800-PC26).
- * The mobility of HIF-1 alpha induced by desferrioxamine or cobalt chloride treatment differs from the mobility of the hypoxia-induced protein. The reason is not known.
- Limitations:** This product is for research use only and is not approved for use in humans or in clinical diagnosis. This product is guaranteed for 6 months from date of receipt.
- Novus Specific References:**
1. Liu, X., et al. Hypoxia preconditioning of cardiomyocytes and cardioprotection: phosphorylation of HIF-1 alpha induced by p42/p44 mitogen-activated protein kinases is involved. *Pathophysiology*. 00: 1-5 (2003).
 2. Julie Litz and Geoffrey W. Krystal. Imatinib inhibits c-Kit-induced hypoxia-inducible factor-1 activity and vascular endothelial growth factor expression in small cell lung cancer cells. *Mol Cancer Ther*. 5: 1415-1422 (2006).
 3. Liao, D., et al. Hypoxia-Inducible Factor-1{alpha} Is a Key Regulator of Metastasis in a Transgenic Model of Cancer Initiation and Progression. *Cancer Research* 67: 563-572 (2007).
 4. Bergeron, M., et al. Role of hypoxia-inducible factor-1 in hypoxia-induced ischemic tolerance in neonatal rat brain. *Ann. Neurol*. 48: 285-296 (2000). (Western blot, rat)
 5. Harris, J. R., et al. Bone Marrow-Derived Cells Home to and Regenerate Retinal Pigment Epithelium after Injury. *Investigative Ophthalmology and Visual Science* 47: 2108-2113 (2006).
 6. Harvey, A. J., et al. Regulation of Gene Expression in Bovine Blastocysts in Response to Oxygen and the Iron Chelator Desferrioxamine. *Biol. Reprod*. 77: 93-101 (2007).
 7. Ietta, F., et al. Dynamic HIF1A Regulation During Human Placental Development. *Biol. Reprod*. 75: 112-121 (2006).
 8. Lin, Q., et al. Differentiation Arrest by Hypoxia. *J. Biol. Chem*. 281: 30678-30683 (2006).
 9. Wang, J., et al. Cytoprotective Effects of Hypoxia against Cisplatin-Induced Tubular Cell Apoptosis: Involvement of Mitochondrial Inhibition and p53 Suppression. *J. Am. Soc. Nephrol*. 17: 1875-1885 (2006).
 10. Young, H. W. J., et al. Central Role of Muc5ac Expression in Mucous Metaplasia and Its Regulation by Conserved 5' Elements. *Am. J. Resp. Cell Molec. Biol*. 37: 273-290 (2007).
 11. Kim J-w, Gao P, Liu Y-C, Semenza GL, Dang CV. Hypoxia-Inducible Factor 1 and Dysregulated c-Myc Cooperatively Induce Vascular Endothelial Growth Factor and Metabolic Switches Hexokinase 2 and Pyruvate Dehydrogenase Kinase 1. *Mol Cell Biol*. 2007 November 1, 2007;27(21):7381-93.
 12. Sikder D, Kodadek T. The neurohormone orexin stimulates hypoxia-inducible factor-1 activity. *Genes & Dev*. 2007 November 15, 2007;21(22):2995-3005.
 13. Singh RP, Gu M, Agarwal R. Silibinin Inhibits Colorectal Cancer Growth by Inhibiting Tumor Cell Proliferation and Angiogenesis. *Cancer Res*. March 15, 2008;68(6):2043-50. (immunohistochemistry)
 14. Lauzier M-C, Robitaille GA, Chan DA, Giaccia AJ, Richard DE. BiPS, a matrix


- metalloprotease inhibitor, is a novel and potent activator of hypoxia-inducible factors. *Mol Pharmacol.* April 18, 2008;mol.108.045690.
15. Ohtomo S, Nangaku M, Izuhara Y, Takizawa S, Strihou CvYd, Miyata T. Cobalt ameliorates renal injury in an obese, hypertensive type 2 diabetes rat model. *Nephrol Dial Transplant.* April 1, 2008;23(4):1166-72.
 16. Piantadosi CA, Suliman HB. Transcriptional Regulation of SDHa Flavoprotein by Nuclear Respiratory Factor-1 Prevents Pseudo-hypoxia in Aerobic Cardiac Cells. *J Biol Chem.* April 18, 2008;283(16):10967-77.
 17. Zhang H, Bosch-Marce M, Shimoda LA, et al. Mitochondrial Autophagy Is an HIF-1-dependent Adaptive Metabolic Response to Hypoxia. *J Biol Chem.* April 18, 2008;283(16):10892-903.
 18. Nikinmaa M, Leveelahti L, Dahl E, et al. Population origin, development and temperature of development affect the amounts of HSP70, HSP90 and the putative hypoxia-inducible factor in the tadpoles of the common frog *Rana temporaria*. *J Exp Biol* 2008;211(12):1999-2004.
 19. Meyuhas R, Pikarsky E, Tavor E, et al. A Key Role for Cyclic AMP-Responsive Element Binding Protein in Hypoxia-Mediated Activation of the Angiogenesis Factor CCN1 (CYR61) in Tumor Cells. *Mol Cancer Res* 2008;6(9):1397-1409.
 20. Trollmann R, Strasser K, Keller S, et al. Placental HIFs as markers of cerebral hypoxic distress in fetal mice. *Am J Physiol Regulatory Integrative Comp Physiol* 2008:00053.02008.
 21. Westfall, S., Sachdev, S., Das, P., et al. Identification of Oxygen-Sensitive Transcriptional Programs in Human Embryonic Stem Cells. *Stem Cells and Development.* Oct. 2008: 869-882.
 22. Land S., and Wilson, S. Redox Regulation of Lung Development and Perinatal Lung Epithelial Function. *Antioxidants & Redox Signaling.* Jan. 2005: 92-107.

General References:


1. Wang, G.L. and Semenza, G.L. Purification and Characterization of Hypoxia-inducible Factor 1. *J. of Biol. Chem.* 270(3): 1230-1237 (1995).
2. Semenza, G.L. Hypoxia-inducible factor 1 and the molecular physiology of oxygen homeostasis. *J. Lab Clinical Medicine.* 131: 207-214 (1998).
3. Kaelin, W.G. Many vessels, faulty gene. *Nature.* 399: 203-204 (1999).
4. Harvey, A.J., et al. Oxygen-Regulated Gene Expression in Bovine Blastocysts. *Biol. Reprod.* 71: 1108-1119 (2004).

Gene Id: 3091
Gene System: HIF1A
Reference Sequence: Q16665

Image(s)



Western Blot analysis of NB 100-134, anti-HIF-1 alpha, on normoxic and hypoxic nuclear rat cell lysates.



Kidney, Renal Tubular Epithelium in Cortex 40x