Product Datasheet

SFRS5 Antibody (2C8) H00006430-M03A

Unit Size: 0.2 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/H00006430-M03A

Updated 6/27/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/H00006430-M03A



H00006430-M03A

SFRS5 Antibody (2C8)


SFRS5 Antibody (2C8)	
Product Information	
Unit Size	0.2 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	2C8
Preservative	No Preservative
Isotype	IgM Kappa
Purity	Unpurified
Buffer	Ascites
Product Description	
Description	Quality control test: Antibody Reactive Against Recombinant Protein.
Host	Mouse
Gene ID	6430
Gene Symbol	SRSF5
Species	Human
Specificity/Sensitivity	SFRS5 - splicing factor, arginine/serine-rich 5 (2C8)
Immunogen	SFRS5 (AAH18823.1, 1 a.a. ~ 272 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa. MSGCRVFIGRLNPAAREKGVERFFKGYGRIRDIDLKRGFGFVEFEDPRDADDA VYELDGKELCSERVTIEHARARSRGGRGRGRYSDRFSSRRPRNDRRNAPPVR TENRLIVENLSSRVSWQDLKDFMRQAGEVTFADAHRPKLNEGVVEFASYGDLK NAIEKLSGKEINGRKIKLIEGSKRHSRSRSRSRSRSRSRSRSRSRSRSRSRSRSRSRSRSR
Notes	This product is produced by and distributed for Abnova, a company based in Taiwan.
Product Application Details	
Applications	Western Blot, ELISA
Recommended Dilutions	Western Blot, ELISA
Application Notes	Antibody reactivity against Recombinant Protein with GST tag on ELISA and



WB. GST tag alone is used as a negative control.

Images

Western Blot: SFRS5 Antibody (2C8) [H00006430-M03A] - Analysis of SFRS5 expression in transfected 293T cell line by SFRS5 monoclonal antibody (M03A), clone 2C8. Lane 1: SFRS5 transfected lysatE (31.3 KDa). Lane 2: Non-transfected lysate.



Publications

Martinez-Pizarro A, Dembic M, Perez B et al. Intronic PAH gene mutations cause a splicing defect by a novel mechanism involving U1snRNP binding downstream of the 5 splice site. PLoS Genet 2018-04-23 [PMID: 29684050]

Bruun GH, Doktor TK, Andresen BS. A synonymous polymorphic variation in ACADM exon 11 affects splicing efficiency and may affect fatty acid oxidation. Molecular Genetics and Metabolism. 2013-01-01 [PMID: 23810226]



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966 novus@novusbio.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/H00006430-M03A

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

