

# Product Datasheet

## PLC/PRF/5 Nuclear Cell Lysate L011V3

Unit Size: 0.05 mg

Store at -80C. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/L011V3](http://www.novusbio.com/L011V3)

Updated 6/27/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/L011V3](http://www.novusbio.com/reviews/destination/L011V3)



**L011V3**

PLC/PRF/5 Nuclear Cell Lysate

**Product Information**

<b>Unit Size</b>	0.05 mg
<b>Concentration</b>	2 mg/ml
<b>Storage</b>	Store at -80C. Avoid freeze-thaw cycles.
<b>Buffer</b>	In ready-to-use 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue).

**Product Description**

<b>Description</b>	Quality control test: 12.5% SDS-PAGE Stained with Coomassie Blue. Nuclear extract cell lysate (denatured)
<b>Species</b>	Human
<b>Specificity/Sensitivity</b>	PLC/PRF/5 (human hepatoma) nuclear extract lysate (denatured)
<b>Preparation Method</b>	Nuclear extract was prepared by using a modified protocol of Dignam et al. Cells were Harvested and homogenized in Buffer A, and then centrifugated at 25,000 g for 20 minutes to remove cytoplasm and pellet the nuclei. The pellet was re-suspended in Buffer C, and then the suspensions were centrifuged to collect nuclear extract. The supernatant was dialyzed against Buffer D. The dialysate was then centrifuged, divided into aliquots, and stored at -80C. The protein concentration was determined by the method of Bradford. The lysate was adjusted to 2.5 mg/ml, and then mixed with 5X Sample Buffer to become final 2 mg/ml in 1X Sample Buffer. The lysate was heated at 95C for 5 min, and cooled rapidly.
<b>Notes</b>	This product is produced by and distributed for Abnova, a company based in Taiwan.
<b>Lysate Type</b>	Cell
<b>Lysate Tissue</b>	Liver
<b>Lysate Protein State</b>	Denatured
<b>Lysate Subcellular Fraction</b>	Nuclear

**Product Application Details**

<b>Applications</b>	Western Blot
<b>Recommended Dilutions</b>	Western Blot
<b>Application Notes</b>	The lysate is ready to load on SDS-PAGE for Western blotting.



## Novus Biologicals USA

10730 E. Briarwood Avenue

Centennial, CO 80112

USA

Phone: 303.730.1950

Toll Free: 1.888.506.6887

Fax: 303.730.1966

[novus@novusbio.com](mailto:novus@novusbio.com)

## Bio-Techne Canada

21 Canmotor Ave

Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400

Toll Free: 855.668.8722

Fax: 905.827.6402

[canada.inquires@bio-techne.com](mailto:canada.inquires@bio-techne.com)

## Bio-Techne Ltd

19 Barton Lane

Abingdon Science Park

Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15

Fax: (44) (0) 1235 533420

[info.EMEA@bio-techne.com](mailto:info.EMEA@bio-techne.com)

## General Contact Information

[www.novusbio.com](http://www.novusbio.com)

Technical Support: [technical@novusbio.com](mailto:technical@novusbio.com)

Orders: [orders@novusbio.com](mailto:orders@novusbio.com)

General: [novus@novusbio.com](mailto:novus@novusbio.com)

## Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis.

Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/L011V3](http://www.novusbio.com/reviews/submit/L011V3)

Earn gift cards/discounts by submitting a publication using this product:

[www.novusbio.com/publications](http://www.novusbio.com/publications)

