

Product Datasheet

PLC/PRF/5 Nuclear Cell Lysate L011V4

Unit Size: 0.05 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/L011V4

Updated 6/27/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/L011V4



L011V4**PLC/PRF/5 Nuclear Cell Lysate****Product Information**

Unit Size	0.05 mg
Concentration	2 mg/ml
Storage	Store at -80C. Avoid freeze-thaw cycles.
Buffer	20mM HEPES pH 7.9, 20%(v/v) glycerol, 50mM KCl, 0.2 mM EDTA, 0.5 mM DTT & 0.5 mM PMSF.

Product Description

Description	Quality control test: 12.5% SDS-PAGE Stained with Coomassie Blue. Nuclear extract cell lysate (non-denatured)
Species	Human
Specificity/Sensitivity	PLC/PRF/5 (human hepatoma) nuclear extract lysate (non-denatured)
Preparation Method	Nuclear extract was prepared by using a modified protocol of Dignam et al. Cells were Harvested and homogenized in Buffer A, and then centrifuged at 25,000 g for 20 minutes to remove cytoplasm and pellet the nuclei. The pellet was re-suspended in Buffer C, and then the suspensions were centrifuged to collect nuclear extract. The supernatant was dialyzed against Buffer D. The dialysate was then centrifuged, divided into aliquots, and stored at -80C. The protein concentration was determined by the method of Bradford. The lysate was adjusted to 2 mg/ml.
Notes	This product is produced by and distributed for Abnova, a company based in Taiwan.
Lysate Type	Cell
Lysate Tissue	Liver
Lysate Protein State	Native
Lysate Subcellular Fraction	Nuclear

Product Application Details

Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot, Immunoprecipitation
Application Notes	Use it directly for immuno-precipitation, or heat lysate with SDS gel loading buffer to 95C for 5 minutes followed by rapid cooling for western blot application. If dissociating conditions are required, add reducing agent prior to heating.



Novus Biologicals USA

10730 E. Briarwood Avenue

Centennial, CO 80112

USA

Phone: 303.730.1950

Toll Free: 1.888.506.6887

Fax: 303.730.1966

novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave

Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400

Toll Free: 855.668.8722

Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane

Abingdon Science Park

Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15

Fax: (44) (0) 1235 533420

info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis.

Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/L011V4

Earn gift cards/discounts by submitting a publication using this product:

www.novusbio.com/publications