

Product Datasheet

Y-79 Nuclear Cell Lysate L042V4

Unit Size: 0.05 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/L042V4

Updated 6/27/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/L042V4



L042V4**Y-79 Nuclear Cell Lysate**

Product Information	
Unit Size	0.05 mg
Concentration	2 mg/ml
Storage	Store at -80C. Avoid freeze-thaw cycles.
Buffer	20mM HEPES pH 7.9, 20%(v/v) glycerol, 50mM KCl, 0.2 mM EDTA, 0.5 mM DTT & 0.5 mM PMSF.
Product Description	
Description	Quality control test: 12.5% SDS-PAGE Stained with Coomassie Blue. Nuclear extract cell lysate (non-denatured)
Species	Human
Specificity/Sensitivity	Y-79 (human retinoblastoma) nuclear extract lysate (non-denatured)
Preparation Method	Nuclear extract was prepared by using a modified protocol of Dignam et al. Cells were Harvested and homogenized in Buffer A, and then centrifugated at 25,000 g for 20 minutes to remove cytoplasm and pellet the nuclei. The pellet was re-suspended in Buffer C, and then the suspensions were centrifuged to collect nuclear extract. The supernatant was dialyzed against Buffer D. The dialysate was then centrifuged, divided into aliquots, and stored at -80C. The protein concentration was determined by the method of Bradford. The lysate was adjusted to 2 mg/ml.
Notes	This product is produced by and distributed for Abnova, a company based in Taiwan.
Lysate Type	Cell
Lysate Tissue	Visual
Lysate Tissue Condition	Retinoblastoma
Lysate Protein State	Native
Lysate Subcellular Fraction	Nuclear
Product Application Details	
Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot, Immunoprecipitation
Application Notes	Use it directly for immuno-precipitation, or heat lysate with SDS gel loading buffer to 95C for 5 minutes followed by rapid cooling for western blot application. If dissociating conditions are required, add reducing agent prior to heating.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/L042V4

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

