## **Product Datasheet**

# RAG1AP1 Antibody NB100-1257

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

**Publications: 1** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-1257

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB100-1257



### NB100-1257

RAG1AP1 Antibody

TO THE TAINING GAY	
Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA
Product Description	
Description	Please note, this antibody is considered Innovators Grade. Innovators Grade antibodies are generally unvalidated and require additional characterization for most new species/applications. Novus has made these antibodies available through our Innovators Reward program. Complete an online review with image, detailing your positive or negative results. In return, you receive a discount voucher for 100% of the purchase price of the reviewed product. Please contact us at innovators@novusbio.com for more details.
Host	Goat
Gene ID	55974
Gene Symbol	SLC50A1
Species	Human
Specificity/Sensitivity	This antibody is expected to recognise the reported isoforms.
Immunogen	Peptide with sequence C-EQDRNYWLLQT corresponding to C-Terminus according to NP_061333.2, NP_001116309.1, NP_001116311.1.

<b>Product Application Details</b>	
Applications	Peptide ELISA
Recommended Dilutions	Peptide ELISA Detection limit 1:8000
Application Notes	WB: Preliminary experiments gave no signal but low background in human liver lysate at up to 1 ug/ml.

## **Publications**

Tagoh H, Kishi H, Muraguchi A. Molecular cloning and characterization of a novel stromal cell-derived cDNA encoding a protein that facilitates gene activation of recombination activating gene (RAG)-1 in human lymphoid progenitors. Biochem Biophys Res Commun 1996-04-25 [PMID: 8630032]





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NB100-1257**

NBL1-15126 RAG1AP1 Overexpression Lysate

HAF017 Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1257

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



