Product Datasheet

SH3BP1 Antibody NB100-1398

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-1398

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB100-1398



NB100-1398


SH3BP1 Antibody

| Unit Size 0.1 mg Concentration 0.5 mg/ml Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Description Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Cacc-2 cells is reported in scientific literature (Elbediwy et al., J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated Mw of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in the interstital parenchyma and of lymphocytes in a glomerulus. | Shabri Antibudy | |
|---|-----------------------------|--|
| Concentration Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Description Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al., J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Product Information | |
| Storage Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Description Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J. Biol. Chem, 2012), WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band aboserved in human lung and kidney lysates (calculated MW of 75.7 kDa band aboserved in lymphoid cells in NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Unit Size | 0.1 mg |
| Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Description Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al., J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney stains the cytoplasm of lymphoid cells in NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Concentration | 0.5 mg/ml |
| Preservative Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Goat Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al., J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Storage | Store at -20C. Avoid freeze-thaw cycles. |
| IgG | Clonality | Polyclonal |
| Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Preservative | 0.02% Sodium Azide |
| Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Isotype | IgG |
| Product Description Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Purity | Immunogen affinity purified |
| Novus Biologicals Goat SH3BP1 Antibody (NB100-1398) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host | Buffer | Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA |
| validated for use in IHC, WB and ELISA. Anti-SH3BP1 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee. Host Goat Gene ID 23616 Gene Symbol SH3BP1 Species Human Immunogen Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Product Description | |
| Gene Symbol SH3BP1 Species Human Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Description | publication. All Novus Biologicals antibodies are covered by our 100% |
| Gene Symbol SH3BP1 Species Human Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Host | Goat |
| Species Human | Gene ID | 23616 |
| Peptide with sequence C-PQPRPRSLASETN corresponding to C-Terminus according to NP_061830.3. Product Application Details | Gene Symbol | SH3BP1 |
| Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Species | Human |
| Applications Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Immunogen | |
| Recommended Dilutions Western Blot 1 - 3 ug/ml, Immunohistochemistry 0.3 ug/ml, Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Product Application Details | |
| Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit 1:16000 Application Notes ICC successful usage in Caco-2 cells is reported in scientific literature (Elbediwy et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Applications | |
| et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in | Recommended Dilutions | Immunohistochemistry-Paraffin 1:10 - 1:500, Peptide ELISA Detection limit |
| | Application Notes | et al, J.Biol. Chem, 2012). WB: Approx. 70 kDa band observed in human lung and kidney lysates (calculated MW of 75.7 kDa band according to NP_061830.3). IHC-P: Human kidney stains the cytoplasm of lymphoid cells in |




Images

Western Blot: SH3BP1 Antibody [NB100-1398] - SH3BP1 in Human Brain lysate using NB100-1398 at 1 ug/ ml.



Immunohistochemistry-Paraffin: SH3BP1 Antibody [NB100-1398] - NB100-1398 (0.3ug/ml) staining of paraffin embedded Human Kidnay. Cytoplasmic staining of lymphocytes A) in the interstitial parenchyma of the kidney. B) in a glomerulus of the kidney. Microwaved antigen retrieval with citrate buffer pH 6, HRP-staining.



Publications

Dunham I, Shimizu N, Roe BA et al. The DNA sequence of human chromosome 22. Nature 1999-12-02 [PMID: 10591208]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB100-1398

NB820-59231 Human Kidney Whole Tissue Lysate (Adult Whole Normal)

NB100-1398PEP SH3BP1 Peptide

HAF017 Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1398

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



