Product Datasheet

BAF53A Antibody NB100-1421

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-1421

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-1421



NB100-1421


BAF53A Antibody

BAF53A Antibody	
Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA
Product Description	
Description	Novus Biologicals Goat BAF53A Antibody (NB100-1421) is a polyclonal antibody validated for use in WB, ELISA, Flow and ICC/IF. Anti-BAF53A Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	86
Gene Symbol	ACTL6A
Species	Human, Mouse, Rat
Specificity/Sensitivity	This antibody is expected to recognise both BAF53A and BAF53B, which are almost identical. Reported variants represent identical protein (NP_817126.1; NP_829888.1).
Immunogen	Peptide with sequence YEEGGKQCVERKCP corresponding to C-Terminus according to NP_004292.1, NP_817126.1, NP_057272.1.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Peptide ELISA
Recommended Dilutions	Western Blot 0.5 - 2 ug/mL, Flow Cytometry 10 ug/mL, Immunocytochemistry/ Immunofluorescence 10 ug/mL, Peptide ELISA Detection limit 1:6000
Images	
Western Blot: BAF53A Antibo	ody [NB100-1421] - Staining of NIH3T3 tein in RIPA buffer). Antibody at 0.5 µg/ml


Images Western Blot: BAF53A Antibody [NB100-1421] - Staining of NIH3T3 nuclear cell lysate (35 ug protein in RIPA buffer). Antibody at 0.5 ug/mL. Detected by chemiluminescence. 250kDa 150kDa 75kDa 25kDa 20kDa 15kDa




Immunocytochemistry/Immunofluorescence: BAF53A Antibody [NB100-1421] - Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).




Flow Cytometry: BAF53A Antibody [NB100-1421] - Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



Western Blot: BAF53A Antibody [NB100-1421] - Staining of Rat Skeletal Muscle lysate (35 ug protein in RIPA buffer). Antibody at 2 ug/mL. Detected by chemiluminescence.



Immunocytochemistry/Immunofluorescence: BAF53A Antibody [NB100-1421] - Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



Publications

Harata M, Mochizuki R, Mizuno S. Two isoforms of a human actin-related protein show nuclear localization and mutually selective expression between brain and other tissues. Biosci Biotechnol Biochem 1999-05-01 [PMID: 10380635]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB100-1421

NB800-PC1 HeLa Whole Cell Lysate

HAF017 Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1421

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



