

Product Datasheet

SKIV2L2 Antibody NB100-1574

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 1 **Publications: 5**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1574

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1574



NB100-1574

SKIV2L2 Antibody


Product Information	
Unit Size	0.1 ml
Concentration	0.2 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	TBS and 0.1% BSA
Target Molecular Weight	118 kDa

Product Description	
Description	Novus Biologicals Rabbit SKIV2L2 Antibody (NB100-1574) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. Anti-SKIV2L2 Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	23517
Gene Symbol	MTREX
Species	Human, Mouse
Immunogen	The immunogen recognized by this antibody maps to a region between residues 100 and 150 of human SuperKiller Viralicidic Activity 2-Like 2 using the numbering given in entry NP_056175.2 (GeneID 23517).


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Immunoprecipitation, Immunocytochemistry, Knockdown Validated
Recommended Dilutions	Western Blot 1:2000-1:10000, Immunohistochemistry 1:100-1:500, Immunoprecipitation 2 - 5 ug/mg lysate, Immunohistochemistry-Paraffin 1:100-1:500, Immunocytochemistry 1:100 - 1:500, Knockdown Validated
Application Notes	For WB: When blotting for mouse SKIV2L2, lower dilutions of antibody and/or longer exposure times may be required. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections. SKIV2L2 antibody validated for WB from a verified customer review.

Images


Western Blot: SKIV2L2 Antibody [NB100-1574] - Analysis of SKIV2L2 in HeLa cell lysates treated with (1) si-control and (2) SKIV2L2 siRNA using anti-SKIV2L2 antibody. Image from verified customer review.




Western Blot: SKIV2L2 Antibody [NB100-1574] - Whole cell lysate (50 ug) from HeLa, 293T, Jurkat, mouse TCMK-1, and mouse NIH3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit antiSKIV2L2 antibody used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



Immunohistochemistry-Paraffin: SKIV2L2 Antibody [NB100-1574] - FFPE section of human lung carcinoma. Antibody: Affinity purified rabbit anti-SKIV2L2 used at a dilution of 1:200 (1ug/ml). Detection: DAB



SKIV2L2 Antibody [NB100-1574] - Detection of human SKIV2L2 by western blot of immunoprecipitates. Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-SKIV2L2 antibody NB100-1574 used for IP at 3 ug per reaction. SKIV2L2 was also immunoprecipitated by rabbit anti-SKIV2L2 antibody NB100-1575. For blotting immunoprecipitated SKIV2L2, NB100-1574 was used at 1 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



SKIV2L2 Antibody [NB100-1574] - Affinity purified rabbit anti-SKIV2L2 used at a dilution of 1:200 (1ug/ml).



Publications

Ogami K, Richard P, Chen Y et al. An Mtr4/ZFC3H1 complex facilitates turnover of unstable nuclear RNAs to prevent their cytoplasmic transport and global translational repression. *Gens & Dev.* 2017-07-21 [PMID: 28733371] (Western Blot, Human)

Chen Y Studies on RNA Regulation: From Enhancer RNAs to RBBP6 Isoform3 Thesis 2019-01-01

Ogami K, Richard P, Chen Y et al. An Mtr4/ZFC3H1 complex facilitates turnover of unstable nuclear RNAs to prevent their cytoplasmic transport and global translational repression *Genes Dev.* 2017 Jul 21 [PMID: 28733371]

Ogami K, Richard P, Chen Y et al. An Mtr4/ZFC3H1 complex facilitates turnover of unstable nuclear RNAs to prevent their cytoplasmic transport and global translational repression. *Gens & Dev.* 2017 Jul 21 [PMID: 28733371]

Nag A, Steitz JA. Tri-snRNP-associated proteins interact with subunits of the TRAMP and nuclear exosome complexes, linking RNA decay and pre-mRNA splicing. *RNA Biol.* 2012-03-01 [PMID: 22336707] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-1574

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1574

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

