Product Datasheet

BTF Antibody NB100-1704

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-1704

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-1704



NB100-1704


BTF Antibody

Product Information 0.1 ml Concentration 0.2 mg/ml Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer TBS and 0.1% BSA Product Description Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.		
Concentration O.2 mg/ml Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative O.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer TBS and 0.1% BSA Product Description Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Product Information	
Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Immunogen affinity purified Buffer TBS and 0.1% BSA Product Description Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen Evaporation Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Unit Size	0.1 ml
Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer TBS and 0.1% BSA Product Description Host Rabbit Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Concentration	0.2 mg/ml
Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer TBS and 0.1% BSA Product Description Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Storage	Store at 4C. Do not freeze.
Isotype IgG Purity Immunogen affinity purified Buffer TBS and 0.1% BSA Product Description Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of Iysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Clonality	Polyclonal
Purity Immunogen affinity purified Buffer TBS and 0.1% BSA Product Description Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Preservative	0.09% Sodium Azide
Buffer TBS and 0.1% BSA Product Description Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Isotype	IgG
Product Description Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Purity	Immunogen affinity purified
Host Rabbit Gene ID 9774 Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Buffer	TBS and 0.1% BSA
Gene Symbol BCLAF1 Species Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Product Description	
Gene Symbol BCLAF1 Human, Mouse Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Host	Rabbit
Species Human, Mouse The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Gene ID	9774
Immunogen The immunogen recognized by this antibody maps to a region between residues 150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Gene Symbol	BCLAF1
150 and 200 of human BCL2-Associated Transcription Factor 1 using the numbering given in entry NP_055554.1 (GeneID 9774). Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Recommended Dilutions Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Species	Human, Mouse
Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation Western Blot 1:2000-1:10000, Immunohistochemistry 1:100 - 1:1000, Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10- 1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Immunogen	150 and 200 of human BCL2-Associated Transcription Factor 1 using the
Immunoprecipitation	Product Application Details	
Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10- 1:500 Application Notes Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Applications	
	Recommended Dilutions	Immunoprecipitation 1 to 4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-
	Application Notes	




Images


Western Blot: BTF Antibody [NB100-1704] - Whole cell lysate from HeLa (5, 15, 50 ug loaded) and 293T cells (T; 50 ug) probed with anti-BTF antibody at 0.04 ug/ml.




Immunohistochemistry: BTF Antibody [NB100-1704] - Sample: FFPE section of human skin carcinoma. Antibody: Affinity purified rabbit anti-BTF used at a dilution of 1:1,000 (0.2ug/ml). Detection: DAB staining using Immunohistochemistry Accessory Kit.



Western Blot: BTF Antibody [NB100-1704] - : Whole cell lysate (50 ug) from TCMK-1, 4T1, and CT26.WT cells.



Immunoprecipitation: BTF Antibody [NB100-1704] - Whole cell lysate from HeLa (5, 15 and 50 ug for WB; 1 mg for IP, 20% of IP loaded) and 293T (T; 50 ug) cells.



Publications

Lee YY, Yu YB, Gunawardena HP et al. BCLAF1 is a radiation-induced H2AX-interacting partner involved in GH2AX-mediated regulation of apoptosis and DNA repair. Cell Death Dis. 2012-07-26 [PMID: 22833098]

Lee SH, Kalejta RF, Kerry J et al. BcIAF1 restriction factor is neutralized by proteasomal degradation and microRNA repression during human cytomegalovirus infection. Proc Natl Acad Sci U S A 2012-06-01 [PMID: 22645331]

Ziegelbauer JM, Sullivan CS, Ganem D et al. Tandem array-based expression screens identify host mRNA targets of virus-encoded microRNAs. Nat Genet 2009-01-01 [PMID: 19098914]

Liu H, Lu ZG, Miki Y et al. Protein kinase C delta induces transcription of the TP53 tumor suppressor gene by controlling death-promoting factor Btf in the apoptotic response to DNA damage. Mol Cell Biol 2007-12-01 [PMID: 17938203]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB100-1704

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP1-89924PEP BTF Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1704

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

