

Product Datasheet

PIP Antibody (23A3) NB100-1898

Unit Size: 0.5 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1898

Updated 3/30/2022 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1898



NB100-1898

PIP Antibody (23A3)

Product Information	
Unit Size	0.5 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	23A3
Preservative	Sodium Azide
Isotype	IgG2a Kappa
Purity	Tissue culture supernatant
Buffer	No buffer
Product Description	
Host	Mouse
Gene ID	5304
Gene Symbol	PIP
Species	Human
Reactivity Notes	Human.
Specificity/Sensitivity	This antibody is specific to a 15 kD monomer protein called Gross Cystic Disease Fluid Protein 15 (GCDFP 15). Gross cystic disease of the breast is benign premenopausal disorder in which cysts are predominant. These cysts appear to be formed from excessive apocrine cystic secretions, one of which is GCDFP-15. This antibody can be used for staining of breast carcinoma, salivary duct carcinoma, and apocrine epithelia.
Immunogen	Recombinant protein encoding the excreted domain of human GCDFP-15.
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry 1:10-1:500, Immunohistochemistry-Paraffin 1:10-1:500
Application Notes	This antibody may be diluted to a titer of 1:10 to 1:20 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user. High temperature antigen unmasking with 10mM citrate buffer, pH 6.0 is required prior to immunostaining. Suggested incubation period of 60 minutes at room temperature.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1898

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

