

Product Datasheet

VEGFR2/KDR/FIk-1 Antibody [Biotin] NB100-2382B

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-2382B

Updated 9/10/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-2382B



NB100-2382B**VEGFR2/KDR/Flk-1 Antibody [Biotin]**

Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Conjugate	Biotin
Purity	Immunogen affinity purified
Buffer	PBS
Product Description	
Host	Rabbit
Gene ID	3791
Gene Symbol	KDR
Species	Human, Rat, Hamster
Reactivity Notes	Human. Predicted to react with mouse based on 100% sequence homology. Immunogen sequence has 79% homology to rat. Hamster reactivity reported in scientific literature (PMID: 23763710). Rat reactivity reported in scientific literature (PMID: 27832737).
Marker	Endothelial Cell Marker
Specificity/Sensitivity	This has only been shown to recognize the native form (180-190 kDa protein) of VEGFR-2, known as Flk-1/KDR/VEGFR 2 (Vascular Endothelial Growth Factor Receptor 2), though it was raised against the precursor form.
Immunogen	A synthetic peptide made to a C-terminal region of the mouse VEGF Receptor 2 protein (between residues 1300-1367). [Swiss-Prot# P35918]
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence
Recommended Dilutions	Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence
Application Notes	<p>This VEGF Receptor 2 antibody is useful for Immunocytochemistry/Immunofluorescence and Western blot, where a doublet is seen at ~150 kDa with CSF-1R/VEGFR2 chimera transfected lysates representing VEGFR-2. A non-specific band is also seen at ~85 kDa. In ICC/IF punctate membrane staining was observed in Hela cells.</p> <p>The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.</p>



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-2382B

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

