Product Datasheet

BcI-2 Antibody NB100-56101

Unit Size: 0.05 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 10

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-56101

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-56101



NB100-56101


Bcl-2 Antibody

BCI-2 Antibody	
Product Information	
Unit Size	0.05 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Unpurified
Buffer	Whole antisera
Product Description	
Host	Rabbit
Gene ID	596
Gene Symbol	BCL2
Species	Human, Mouse, Rat, Canine, Gerbil
Immunogen	A synthetic peptide corresponding to amino acids 67-86 (ARTSPLRPLVATAGPALSPV) of mouse and rat Bcl-2 was used as immunogen; GenBank no gb/AAP47159.1. This sequence is 100% conserved in mouse and rat Bcl-2 alpha (236 amino acids) and beta (100 amino acids) isoforms.
Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000-1:2000, Immunohistochemistry, Immunoprecipitation 1:50-

Images

Western Blot: Bcl-2 Antibody [NB100-56101] - Analysis of Bcl-2 in 293T cell lysate (30ug) using anti-Bcl-2 antibody. Image from verified customer review.

Frozen 1:10-1:500



1:200, Immunohistochemistry-Paraffin 1:1000-1:5000, Immunohistochemistry-

Publications

Wu Z, Hu J, Li Y et al. Assessment of renal pathophysiological processes and protective effect of quercetin on contrast-induced acute kidney injury in type 1 diabetic mice using diffusion tensor imaging Redox Report : Communications in Free Radical Research 2024-09-16 [PMID: 39284588]

Shoma Ishikawa, Fuyuki Ishikawa Proteostasis failure and cellular senescence in long ☐ term cultured postmitotic rat neurons Aging Cell 2019-11-25 [PMID: 31762159]

Debarpan Mitra, Depanwita Saha, Gaurav Das, Rimi Mukherjee, Samir Banerjee, Neyaz Alam, Saunak Mitra Mustafi, Partha Nath, Anuj Majumder, Biswanath Majumder, Nabendu Murmu Lupeol synergizes with 5-fluorouracil to combat c-MET/EphA2 mediated chemoresistance in triple negative breast cancer iScience 2023-11-04 [PMID: 38047085]

Polat ME, Sar? E, Tanriverdi LH et al. A potential hepatoprotective effect of ranolazine against methotrexate-induced liver injury in rats Authorea 2023-01-01 (IHC, Rat)

Arman T, Baron J, Lynch K Et al. MCLR-elicited hepatic fibrosis and carcinogenic gene expression changes persist in rats with diet-induced nonalcoholic steatohepatitis through a 4-week recovery period Toxicology 2021-11-01 [PMID: 34740672] (WB, Rat)

Arman T, Lynch KD, Goedken M, Clarke JD Sub-chronic microcystin-LR renal toxicity in rats fed a high fat/high cholesterol diet Chemosphere 2020-10-27 [PMID: 33143886] (IHC-P, Rat)

Sulaiman D, Li J, Devarajan A et al. Paraoxonase 2 protects against acute myocardial ischemia-reperfusion injury by modulating mitochondrial function and oxidative stress via the PI3K/Akt/GSK-3b RISK pathway J. Mol. Cell. Cardiol. 2019-02-23 [PMID: 30802459] (IHC-P, Mouse)

Samdanci ET, Huz M, Ozhan O et al. Cytoprotective effects of molsidomine against methotrexate-induced hepatotoxicity: an experimental rat study Drug Des Devel Ther 2019-12-20 [PMID: 30587924] (IF/IHC, Rat)

Rajamani U, Essop MF. Hyperglycemia-mediated activation of the hexosamine biosynthetic pathway results in myocardial apoptosis. Am J Physiol Cell Physiol. 2010-07-01 [PMID: 20410435]

Krajewska M, Rosenthal RE, Mikolajczyk J et al. Early processing of Bid and caspase-6, -8, -10, -14 in the canine brain during cardiac arrest and resuscitation. Exp Neurol. 2004-10-01 [PMID: 15380478] (IHC-P, WB, Mouse, Canine, Rat)

Details:

Antibodies cited: 1. Caspase-9 (Active/Cleaved), IMG-5705: WB: Fig 1A (recombinant human caspase-9), Fig 4A (recombinant human caspase-9, rat brain), Fig 4B-F (isolated mitochondria from rat liver or heart). Fig 5 (PC12 cells) IHC (P): Fig 2d-k (rat kidne





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB100-56101

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-34889-100ug Recombinant Human Bcl-2 Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56101

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

