

Product Datasheet

BIK Antibody NB100-56109

Unit Size: 0.05 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56109

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56109



NB100-56109


BIK Antibody

Product Information	
Unit Size	0.05 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Unpurified
Buffer	Whole antisera
Product Description	
Host	Rabbit
Gene ID	638
Gene Symbol	BIK
Species	Human, Mouse, Rat
Immunogen	Recombinant full-length human Bik protein was used as immunogen.
Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000-1:2000, Immunohistochemistry, Immunoprecipitation 1:50-1:200, Immunohistochemistry-Paraffin 1:1000-1:5000




Images

Immunohistochemistry-Paraffin: BIK Antibody [NB100-56109] - Tissue section of human Non-Hodgkin's lymphoma (NHL) stained for Bik expression using this antibody at 1:2000. Hematoxylin-eosin counterstain.



Immunohistochemistry-Paraffin: BIK Antibody [NB100-56109] - Tissue section of human kidney stained for Bik expression using this antibody at 1:2000. A high level of Bik signal was seen in the distal collecting tubules, the glomeruli are primarily negative. Hematoxylin-eosin counterstain.



Publications

Fu D, Li P, Sheng Q, Lv Z beta-arrestin-2 enhances intestinal epithelial apoptosis in necrotizing enterocolitis Aging 2019-10-14 [PMID: 31612867] (WB, Mouse, Human)

Suzuki K, Yoshida K, Ueha T et al. Methotrexate upregulates circadian transcriptional factors PAR bZIP to induce apoptosis on rheumatoid arthritis synovial fibroblasts. Arthritis Res. Ther. 2018-03-22 [PMID: 29566767] (WB, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-56109

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
H00000638-P01-10ug	Recombinant Human BIK GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56109

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

