

# Product Datasheet

## HDAC1 Antibody - BSA Free

### NB100-56340

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 2 Publications: 19**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB100-56340](http://www.novusbio.com/NB100-56340)

Updated 4/13/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB100-56340](http://www.novusbio.com/reviews/destination/NB100-56340)



**NB100-56340**

HDAC1 Antibody - BSA Free

**Product Information**

<b>Unit Size</b>	0.1 mg
<b>Concentration</b>	1.0 mg/ml
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Polyclonal
<b>Preservative</b>	0.02% Sodium Azide
<b>Isotype</b>	IgG
<b>Purity</b>	Protein G purified
<b>Buffer</b>	PBS

**Product Description**

<b>Host</b>	Rabbit
<b>Gene ID</b>	3065
<b>Gene Symbol</b>	HDAC1
<b>Species</b>	Human, Mouse, Rat
<b>Reactivity Notes</b>	Mouse reactivity reported in scientific literature (PMID: 25648995). Rat reactivity reported in scientific literature (PMID: 26551542)
<b>Immunogen</b>	This antibody was generated by immunizing rabbits with a mixture of synthetic peptides corresponding to amino acids 1-5, 433-448 and 467-482 of human HDAC1 (Genbank Accession no. Q13547).


**Product Application Details**

<b>Applications</b>	Western Blot, Simple Western, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Knockout Validated
<b>Recommended Dilutions</b>	Western Blot 2 ug/ml, Simple Western 1:200, Immunohistochemistry 1:500. Use reported in multiple pieces of scientific literature, Immunohistochemistry-Paraffin 1:10-1:500. Use reported in scientific literature (PMID 25648995), Immunohistochemistry-Frozen 1:10-1:500. Use reported in scientific literature (PMID 26551542), Knockout Validated
<b>Application Notes</b>	In 293, a 60 kDa band is observed. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. See <a href="#">Simple Western Antibody Database</a> for Simple Western validation: Tested in Hek293 lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:100, apparent MW was 68 kDa




## Images


Western Blot: HDAC1 Antibody [NB100-56340] - Western blot shows lysates of HeLa human cervical epithelial carcinoma parental cell line and HDAC1 knockout (KO) HeLa cell line. PVDF membrane was probed with 2.0 ug/ml of Rabbit Anti-Human HDAC1 Polyclonal Antibody (Catalog # NB100-56340) followed by HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog #HAF008). Specific band was detected for HDAC1 at approximately 65 kDa (as indicated) in the parental HeLa cell line, but is not detectable in the knockout HeLa cell line. This experiment was conducted under reducing conditions.




Western Blot: HDAC1 Antibody [NB100-56340] - Analysis of HDAC1 in HEK293 cell lysate with this antibody.



Immunohistochemistry-Paraffin: HDAC1 Antibody [NB100-56340] - Human testis using HDAC1 antibody at 1:250 on a Bond Rx autostainer (Leica Biosystems). The assay involved 20 minutes of heat induced antigen retrieval (HIER) using 10mM sodium citrate buffer (pH 6.0) and endogenous peroxidase quenching with peroxide block. The sections were incubated with primary antibody for 30 minutes and Bond Polymer Refine Detection (Leica Biosystems) with DAB was used for signal development followed by counterstaining with hematoxylin. Whole slide scanning and capturing of representative images was performed using Aperio AT2 (Leica Biosystems). Staining was performed by Histowiz.



Simple Western: HDAC1 Antibody [NB100-56340] - Simple Western lane view shows a specific band for HDAC1 in 0.5 mg/ml of HEK293 lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



## Publications

Alvarez MEG, McGuire BC, Keating AF Obesity alters the ovarian proteomic response to zearalenone exposure  
Biology of reproduction 2021-04-14 [PMID: 33855340]

Laban H, Sigmund S, Schlereth K et al. NFAT5-dependent transcriptional stress control of endothelial cells prevents maladaptive remodeling of pulmonary arterioles in the hypoxic lung bioRxiv 2023-10-22 (Simple Western, Mouse)

Details:  
Dilution 1:10

Laban H, Siegmund S, Zappe M et al. NFAT5/TonEBP Limits Pulmonary Vascular Resistance in the Hypoxic Lung by Controlling Mitochondrial Reactive Oxygen Species Generation in Arterial Smooth Muscle Cells Cells 2021-11-24 [PMID: 34943801]

Opichka M Placental and Syncytiotrophoblast-Specific GAQ Signaling in the Pathogenesis of Preeclampsia Thesis 2023-01-01 (Simple Western)

Kappert L, Ruzicka P, Kutikhin A Et al. Loss of Nfat5 promotes lipid accumulation in vascular smooth muscle cells FASEB journal : official publication of the Federation of American Societies for Experimental Biology 2021-09-01 [PMID: 34383982] (WB, Mouse)

Lazarchuk P, Hernandez-Villanueva J, Pavlova Mn Et Al. Mutual balance of histone deacetylases HDAC1, HDAC2, and the acetyl reader ATAD2 regulates the level of acetylation of histone H4 on nascent chromatin of human cells Mol. Cell. Biol. 2020-02-03 [PMID: 32015101] (WB, KD, ICC/IF)

Johnson AM The Role of the Tumor Microenvironment on the Progression of Lung Cancer Thesis 2020-01-01 (WB, Mouse, Human)

Zhang, Q;Chao, TC;Patil, VS;Qin, Y;Tiwari, SK;Chiou, J;Dobin, A;Tsai, CM;Li, Z;Dang, J;Gupta, S;Urdahl, K;Nizet, V;Gingeras, TR;Gaulton, KJ;Rana, TM; The long noncoding RNA ROCK1 regulates inflammatory gene expression EMBO J. 2019-04-15 [PMID: 30918008] (WB, Human)

Vlaming H, McLean C, Korthout T, Alemdehy MF. Evolutionarily-conserved chromatin crosstalk generates a DOT1L-dose dependency in thymic lymphoma caused by loss of HDAC1 bioRxiv 2019-01-10 (WB, Mouse)

Wijdeven RH, van Luijn MM, Wierenga-Wolf AF et al. Chemical and genetic control of IFN $\gamma$ -induced MHCII expression. EMBO Rep. 2018-07-18 [PMID: 30021835] (WB, Human)

Laumet G, Garriga J, Chen SR et al. G9a is essential for epigenetic silencing of K(+) channel genes in acute-to-chronic pain transition. Nat. Neurosci. 2015-12-01 [PMID: 26551542] (WB, IHC-Fr, Rat)

Matthias Patrick. Too much or too little, how much HDAC activity is good for you? Blood. 2013-03-14 [PMID: 23493769] (WB)

More publications at <http://www.novusbio.com/NB100-56340>



## Procedures

### Western Blot Protocol for HDAC1 Antibody (NB100-56340)

#### Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 10-25 ug of total protein per lane.
2. Transfer proteins to PVDF membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
3. Stain the membrane with Ponceau S (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
4. Rinse the blot TBS -0.05% Tween 20 (TBST).
5. Block the membrane in 5% Non-fat milk in TBST (blocking buffer) for at least 1 hour.
6. Wash the membrane in TBST three times for 10 minutes each.
7. Dilute primary antibody in 2% Non-fat milk in TBST and incubate overnight at 4C with gentle rocking.
8. Wash the membrane in TBST three times for 10 minutes each.
9. Incubate the membrane in diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) for 1 hour at room temperature.
10. Wash the blot in TBST three times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturer's instructions.

### Immunohistochemistry-Paraffin Protocol for HDAC1 Antibody (NB100-56340)

#### Immunohistochemistry-Paraffin Embedded Sections

##### Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes (keep slides in the sodium citrate buffer at all times).

##### Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in PBS for 5 minutes.
3. Block each section with 100-400 ul blocking solution (1% BSA in PBS) for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul HRP polymer conjugated secondary antibody. Incubate 30 minutes at room temperature.
7. Wash sections three times in wash buffer for 5 minutes each.
8. Add 100-400 ul DAB substrate to each section and monitor staining closely.
9. As soon as the sections develop, immerse slides in deionized water.
10. Counterstain sections in hematoxylin.
11. Wash sections in deionized water two times for 5 minutes each.
12. Dehydrate sections.
13. Mount coverslips.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

[www.novusbio.com](http://www.novusbio.com)  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB100-56340**

NB800-PC6	293 Whole Cell Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB100-56340](http://www.novusbio.com/reviews/submit/NB100-56340)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

