

Product Datasheet

Exportin 4 Antibody - BSA Free NB100-56495

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56495

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56495



NB100-56495

Exportin 4 Antibody - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein G purified
Buffer	PBS containing 0.2% gelatin

Product Description	
Host	Rabbit
Gene ID	64328
Gene Symbol	XPO4
Species	Human, Mouse, Rat, Canine, Chicken, Primate
Reactivity Notes	The amino acid sequence used as immunogen is 100% homologous in human (isoforms CRA_a, _b, _c), rhesus monkey, chimp (isoforms 1-6), 94% homologous in bovine and xenopus, and 88% homologous in zebrafish.
Immunogen	A portion of amino acids 850-950 of human Exportin-4 was used as the immunogen.


Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1-3 ug/ml, Immunohistochemistry, Immunohistochemistry-Paraffin

Images


Western Blot: Exportin 4 Antibody [NB100-56495] - Analysis of Exportin-4 in Jurkat cell lysate in the 1) absence and 2) presence of immunizing peptide using this antibody.




Immunohistochemistry-Paraffin: Exportin 4 Antibody [NB100-56495] - Formalin-fixed, paraffin-embedded human hepatocellular carcinoma stained with TEM8 antibody (5 ug/ml), peroxidase-conjugate and DAB chromogen. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM sodium citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.



Immunohistochemistry-Paraffin: Exportin 4 Antibody [NB100-56495] - Formalin-fixed, paraffin-embedded human liver stained with TEM8 antibody (5 ug/ml), peroxidase-conjugate and DAB chromogen. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM sodium citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.



Immunoprecipitation: Exportin 4 Antibody [NB100-56495] - RASSF1A binds to XPO6 though its SARAH domain. Co-immunoprecipitation of endogenous RASSF1A with endogenous RAN, CRM1/XPO1, XPO4, XPO5, XPO6 & XPO7 from HeLa cell lysates, compared with the IgG control. Upper: graphical representation of the domain structure of full-length RASSF1A & mutant constructs used for mapping RASSF1A/XPO6 interaction. Different domains are abbreviated as follows: SARAH, Salvador-RASSF-Hippo domain; C1, N-terminal C1 type zinc fingers; & RA, Ras-binding domain. Lower: Western blot analysis of MYC immunoprecipitation from the indicated inputs from HeLa cells. Co-immunoprecipitation of endogenous XPO6 with RAN in siRASSF1A from HeLa cell lysates. Co-immunoprecipitation of endogenous XPO6 with endogenous RAN, RASSF1A & MST2 in siRASSF1A & siMST2 HeLa cells. Quantification of the interaction of XPO6 with RASSF1A, RAN & MST2 relative to XPO6 is shown. Error bars derive from three independent experiments & represent the SEM. Co-immunoprecipitation of endogenous RASSF1A with endogenous XPO6 & RAN in siRNA-mediated knockdown of MST2 HeLa cells. Quantification of the interaction of RASSF1A with XPO6, RAN & MST2 relative to RASSF1A is shown. Error bars derive from three independent experiments & represent the SEM. Data information: Two-tailed Student's t-test was used for statistical analysis. *P < 0.05, **P < 0.01. Source data are available online for this figure. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31414556>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Chatzifrangkeskou, M, Pefani, D E Et al. RASSF1A is required for the maintenance of nuclear actin levels. EMBO J 2019-08-15 [PMID: 31414556] (WB, Mouse)

Metwally M, Bayoumi A, Khan A Et al. Copy number variation and expression of exportin-4 associates with severity of fibrosis in metabolic associated fatty liver disease EBioMedicine 2021-08-01 [PMID: 34388518] (WB, Human)

Chatzifrangkeskou M, Pefani D E et al. RASSF1A is required for the maintenance of nuclear actin levels. EMBO J 2019-07-06 [PMID: 31175105] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-56495

NB800-PC2	Jurkat Whole Cell Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56495

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

