

Product Datasheet

TAF15 Antibody - BSA Free

NB100-566

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 9

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-566

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-566



NB100-566

TAF15 Antibody - BSA Free

Product Information

Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)

Product Description

Host	Rabbit
Gene ID	8148
Gene Symbol	TAF15
Species	Human
Specificity/Sensitivity	NB100-566 is specific for human TAFII-68 protein.
Immunogen	A synthetic peptide which maps to a region between residues 175 and 225 of human TAFII68 using the numbering given in Swiss-Prot entry Q92804 (GeneID 8148).


Product Application Details

Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:5000-1:20000, Immunohistochemistry 1:1000 - 1:5000, Immunoprecipitation 1-4 ug/mg lysate, Immunohistochemistry-Paraffin 1:1000 - 1:5000
Application Notes	Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. In some cases, the antibody may be diluted further than indicated. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.




Images

Immunohistochemistry-Paraffin: TAF15 Antibody [NB100-566] - Sample: FFPE section of human testicular seminoma. Antibody: Affinity purified rabbit anti- TAFII68 used at a dilution of 1:5,000 (0.2ug/ml). Detection: DAB



Immunoprecipitation: TAF15 Antibody [NB100-566] - Detection of human TAFII68 on HeLa whole cell lysate using NB100-566. TAFII68 was IPed with rabbit anti-TAFII68 antibodies NB100-566, NB100-567, NB100-568, and another using each at 0.3 ug/mg lysate.



Publications

Duarte Ai, Candeias E, Alves In et Al. Liraglutide Protects Against Brain Amyloid-beta 1-42 Accumulation in Female Mice with Early Alzheimer's Disease-Like Pathology by Partially Rescuing Oxidative/Nitrosative Stress and Inflammation Int J Mol Sci 2020-03-04 [PMID: 32143329]

Fernandes T, Resende R, Silva D Et Al. Structural and Functional Alterations in Mitochondria-Associated Membranes (MAMs) and in Mitochondria Activate Stress Response Mechanisms in an In Vitro Model of Alzheimer s Disease Biomedicines 2021-07-24 [PMID: 34440085]

Gao QY, Zhang HF, Tao J et al. Mitochondrial Fission and Mitophagy Reciprocally Orchestrate Cardiac Fibroblasts Activation Frontiers in Cell and Developmental Biology 2021-01-21 [PMID: 33585469]

Vijayan M, Reddy PH Reduced VDAC1, Maintained Mitochondrial Dynamics and Enhanced Mitochondrial Biogenesis in a Transgenic Tau Mouse Model of Alzheimer's Disease International journal of molecular sciences 2022-08-02 [PMID: 35955694]

Details:

WB dilution used 1:1000, IHC dilution used 1:102

Wongkitikamjorn W, Hosomichi J, Wada E et al. Gestational Intermittent Hypoxia Induces Mitochondrial Impairment in the Geniohyoid Muscle of Offspring Rats Cureus 2022-05-17 [PMID: 35600069]

Marques A, Resende R, Silva D et al. Mitochondrial Alterations in Fibroblasts of Early Stage Bipolar Disorder Patients Biomedicines 2021-05-07 [PMID: 34066918]

Cristina Carvalho, Sónia C. Correia, Raquel Seíça, Paula I. Moreira WWOX inhibition by Zfra1-31 restores mitochondrial homeostasis and viability of neuronal cells exposed to high glucose Cellular and Molecular Life Sciences: CMLS 2022-09-01 [PMID: 35984507]

Kim GH, Kwon I. Distinct roles of hnRNPH1 low-complexity domains in splicing and transcription Proceedings of the National Academy of Sciences 2021-12-14 [PMID: 34873036] (Immunoprecipitation, Western Blot)

Kim E, Kwon I Phase transition of fibrillarin LC domain regulates localization and protein interaction of fibrillarin The Biochemical journal 2021-02-26 [PMID: 33522570]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-566

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-22848	Recombinant Human TAF15 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-566

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

