

Product Datasheet

NFATC1/NFAT2 Antibody - BSA Free NB100-56732

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56732

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56732



NB100-56732


NFATC1/NFAT2 Antibody - BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein G purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Rabbit NFATC1/NFAT2 Antibody - BSA Free (NB100-56732) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and Simple Western. Anti-NFATC1/NFAT2 Antibody: Cited in 5 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	4772
Gene Symbol	NFATC1
Species	Human, Mouse
Specificity/Sensitivity	The amino acid sequence used is 100% identical in human NFATc1 A, B, C and E isoforms. The reported amino acid size of NFATc1 isoforms are as follows: A - 716 aa, B-825 aa, C-930 aa, and E-812 aa.
Immunogen	Amino acids 264-282 (NKRKYSNLNGRQPPYSPHHS) of human NFATc1 were used as immunogen for this antibody.
Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1-3 ug/ml, Simple Western 1:50, Immunohistochemistry reported in scientific literature (PMID 35167499), Immunocytochemistry/ Immunofluorescence 1:20
Application Notes	<p>Immunoreactive bands can be blocked by preincubation of NB100-56732 with immunizing peptide. The NFATc1 antibody may detect multiple isoforms of different sizes depending upon testing material.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See Simple Western Antibody Database for Simple Western validation: Tested in RAW lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:50, apparent MW was 43 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>




Images


Western Blot: NFATC1/NFAT2 Antibody [NB100-56732] - Immortalized mouse podocytes. Image from verified customer review.




Immunocytochemistry/Immunofluorescence: NFATC1/NFAT2 Antibody [NB100-56732] - NFAT2 antibody was tested in Raw 246.7 cells with DyLight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and DyLight 550 (red). Image objective 40x. An antibody dilution of 1:10 was used.



Western Blot: NFATC1/NFAT2 Antibody [NB100-56732] - Analysis of human (A) Ramos and (B) mouse RAW lysate using NFATc1 antibody at 1 ug/ml.



Western Blot: NFATC1/NFAT2 Antibody [NB100-56732] - Analysis of NFATc1 in Ramos cell lysate. (1) Without blocking peptide. (2) With blocking peptide.



Simple Western: NFATC1/NFAT2 Antibody [NB100-56732] - Lane view shows a specific band for NFAT2 in 0.5 mg/ml of RAW lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Publications

Janardhan HP, Dresser K, Hutchinson L, Trivedi CM. Pathological MAPK activation-mediated lymphatic basement membrane disruption causes lymphangiectasia that is treatable with raxoxertinib JCI Insight 2022-09-08 [PMID: 36073544] (Western Blot, Mouse)

Sarna NS, Desai SH, Kaufman BG, Curry NM et Al. Enhanced and sustained T cell activation in response to fluid shear stress iScience 2024-06-17 [PMID: 38883838]

Pham TX, Lee J, Guan J et al. Transcriptional analysis of lung fibroblasts identifies PIM1 signaling as a driver of aging-associated persistent fibrosis JCI insight 2022-02-15 [PMID: 35167499] (IF/IHC, Human)

Song J, Jing ZZ, Hu W et al. alpha-Linolenic Acid Inhibits Receptor Activator of NF-κB Ligand Induced (RANKL-Induced) Osteoclastogenesis and Prevents Inflammatory Bone Loss via Downregulation of Nuclear Factor-KappaB-Inducible Nitric Oxide Synthases (NF-κB-iNOS) Signaling Pathways. Med Sci Monit. 2017-10-24 [PMID: 29061958] (WB, Mouse)

Mack JJ, Mosqueiro TS, Archer BJ et al. NOTCH1 is a mechanosensor in adult arteries Nat Commun. 2017-11-20 [PMID: 29158473] (ICC/IF, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-56732

NBL1-13608	NFATC1/NFAT2 Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56732

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

