Product Datasheet

FoxP1 Antibody (JC12) - BSA Free NB100-65125

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 8

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-65125

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB100-65125



NB100-65125

FoxP1 Antibody (JC12) - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	JC12
Preservative	0.05% Sodium Azide
Isotype	IgG2a
Purity	Protein G purified
Buffer	PBS
Product Description	
Host	Mouse
Gene ID	27086
Gene Symbol	FOXP1
Species	Human, Mouse
Specificity/Sensitivity	Does not recognize closely related molecules FOXP2, FOXP3 or FOXP4.
Immunogen	Human FOXP1 [Uniprot# Q9H334]
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry- Paraffin, Immunoprecipitation, CyTOF-ready
Recommended Dilutions	Western Blot 2ug/ml, Flow Cytometry 1 ug per million cells, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence reported in scientific literature (PMID 15161711), Immunoprecipitation reported in scientific literature (PMID 17586580), Immunohistochemistry-Paraffin 1:200- 1:500, Immunohistochemistry-Frozen 1:10-1:500, CyTOF-ready
Application Notes	Heat induced antigen retrieval with Sodium Citrate buffer pH 6.0 is recommended when using this antibody for IHC-P. This antibody is CyTOF ready.

Images

Western Blot: FOXP1 Antibody (JC12) [NB100-65125] - Western blot analysis of resonicated MCF7 cell lysate (A) and MCF7 cell lysate (B) using FOXP1 antibody at 2 ug/ml.





Immunocytochemistry/Immunofluorescence: FOXP1 Antibody (JC12) [NB100-65125] - FOXP1 antibody was tested at 1:25 in MCF-7 cells with Dylight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and Dylight 550 (red). Image objective 40x.

Immunohistochemistry-Paraffin: FOXP1 Antibody (JC12) [NB100-65125] - IHC analysis of formalin-fixed paraffin-embedded tissue section of malignant stromal tumor of the human small bowel using mouse monoclonal FOXP1 antibody (clone JC12) at 5 ug/ml concentration. The carcinoma cells developed an expected and specific strong nuclear with mild cytoplasmic immunopositivity for FOXP1 protein.

Flow Cytometry: FOXP1 Antibody (JC12) [NB100-65125] - Intracellular flow cytometric staining of 1 x 10^6 CHO (A) and HeLa (B) cells using FOXP1 antibody (dark blue). Isotype control shown in orange. An antibody concentration of 1 ug/1x10^6 cells was used.

Immunohistochemistry-Paraffin: FOXP1 Antibody (JC12) [NB100-65125] - IHC analysis of formalin-fixed paraffin-embedded tissue section of human normal breast using FOXP1 antibody (clone JC12) at 5 ug/ml concentration. The breast ductal/acinar epithelial cells and the myoepithelial cells developed a strong nuclear along with moderate cytoplasmic immuno-positivity for FOXP1 protein.

www.novusbio.com









Immunohistochemistry-Paraffin: FOXP1 Antibody (JC12) [NB100-65125] IHC analysis of formalin-fixed paraffin-embedded tissue section of human normal brain using mouse monoclonal FOXP1 antibody (clone JC12) at 5 ug/ml concentration. The cells in the brain tissue depicted strong specific nuclear along with relatively weak cytoplasmic immunopositivity for FOXP1 protein.

- IHC analysis of formalin-fixed paraffin-embedded tissue section of human normal colon using mouse monoclonal FOXP1 antibody (clone JC12) at 5 ug/ml concentration. Most of the cells depicted an expected

strong nuclear with mild cytoplasmic staining.



Publications

Gars E, Butzmann A, Ohgami R et al. The life and death of the germinal center Ann Diagn Pathol 2019-11-13 [PMID: 31751845] (IF/IHC, Human)

Li B, Samanta A, Song X et al. FOXP3 is a homo-oligomer and a component of a supramolecular regulatory complex disabled in the human XLAAD/IPEX autoimmune disease. Int Immunol. 2007-07-01 [PMID: 17586580] (IP, Human)

Brown PJ, Ashe SL, Leich E et al. Potentially oncogenic B-cell activation-induced smaller isoforms of FOXP1 are highly expressed in the activated B cell-like subtype of DLBCL. Blood. 2008-03-01 [PMID: 18077790] (WB, IF/IHC, Human)

Fox SB, Brown P, Han C et al. Expression of the forkhead transcription factor FOXP1 is associated with estrogen receptor alpha and improved survival in primary human breast carcinomas. Clin Cancer Res, 2004-05-15 [PMID: 15161711] (IF/IHC, WB, ICC/IF, Human)

Loddenkemper C, Maul J, Berg E et al. Analysis of FOXP3 protein expression in human CD4(+)CD25(+) regulatory T cells at the single-cell level. Eur J Immunol. 2006-01-01 [PMID: 16380960] (IF/IHC, Human)

Xie Y, Bulbul MA, Ji L et al. p53 Expression Is a Strong Marker of Inferior Survival in De Novo Diffuse Large B-Cell Lymphoma and May Have Enhanced Negative Effect With MYC Coexpression: A Single Institutional Clinicopathologic Study. Am. J. Clin. Pathol. 2014-04-01 [PMID: 24619762] (IHC-P, Human)

Banham, AH et al. Expression of the FOXP1 transcription factor is strongly associated with inferior survival in patients with diffuse large B-cell lymphoma. Clin Cancer Res11:1065 - 1072. 2005-01-01 [PMID: 15709173]

Banham, AH et al. The FOXP1 winged helix transcription factor is a novel candidate tumor suppressor gene on chromosome 3p. Cancer Res 61:8820 - 8829. 2001-01-01 [PMID: 11751404]



Procedures

Immunohistochemistry-Paraffin protocol for FoxP1 Antibody (NB100-65125)

1. Deparaffinize the tissue sections by immersing the slides in Xylene with two changes for 10 min each. Sections should not get dried at any stage from this point.

- 2. Rehydrate the tissue sections by immersing the slides in decreasing grades of ethanol as follows:
- a. Immerse in 100% ethanol with 2 changes for 5 minutes each
- b. Immerse in 95% ethanol with 2 changes for 5 minutes each
- c. Immerse in 90% ethanol for 5 minutes
- d. Immerse in 70% ethanol for 5 minutes
- e. Immerse in 50% ethanol for 5 minutes
- f. Immerse in distilled water for 5 minutes
- 3. Antigen Retrieval (Microwave Method):

a. Immerse the slides in a microwave compatible tray containing 10 mM Sodium Citrate buffer (pH 6.0) with 0.05% Tween 20.

b. Boil the slides and maintain the sub-boiling temperature for 5 minutes in the microwave. Thereafter, take out the tray very carefully and cool it at room temperature (RT) for about 30 minutes.

c. Wash the slides 3 times, 3 minutes each by immersing them in TBST (Tris Buffered Saline having 0.05% Tween 20).

- 4. Quenching of Endogenous Peroxidase:
- a. Incubate the slides in 3% hydrogen peroxide prepared in methanol for 15 minutes (at RT, in dark conditions).
- b. Wash the slides in TBST 3 times, 3 minutes each.
- 5. Protein Blocking:
- a. Incubate the sections with background sniper solution at RT for 15 minutes (Biocare Medicals, USA).
- b. Wash the sections 3 times, 3 min each by immersing the slides in TBST.
- 6. Primary Antibody:
- a. Dilute the primary antibody at 2-5ug/ml concentration using PBS as a diluent.
- b. Incubate the sections with diluted primary antibody for 90 minutes at RT in a humidified chamber.
- c. Thereafter, wash the slides 4 times, 5 minutes each with TBST.
- 7. Probe (Secondary Reagent):
- a. Incubate with MACH 1 Mouse probe for 15 minutes at RT.
- b. Incubate for 30 min at room temperature with HRP-Polymer (Biocare Medical, USA).
- c. Wash the slides with TBST 4 times, 5 minutes each
- 8. Chromogen:
- a. Mix 32ul of DAB Chromogen with 1 ml of DAB substrate buffer (Biocare Medical, USA).
- a. Apply 200ul DAB mixture/section and incubate at RT in dark conditions (few seconds 5 minutes).
- b. As soon as an appropriate color develops, rinse the slides with deionized water (2-3 brief rinses).
- 9. Counter stain:
- a. Counter stain with Hematoxylin for 30 seconds (Vector Labs, USA).
- b. Wash in deionized water for 1-2 minutes to clear the extra stain.
- c. Incubate the slides in bluing solution or Scott's water twice for 2 minutes each time.
- 10. Dehydrate the sections in increasing grades of alcohols:
- a. 50% alcohol for 1 minute
- b. 70% for 1 minute
- c. 90% for 1 minute
- d. 95% for 1 minute
- e. 100% for 1 minute
- f. Xylene with 2 changes for 2 minutes each

11. Mount with DPX mount and cover-slip glass (Fisher Scientific, USA), carefully not allowing any air bubbles to enter.

NOTE:- This protocol is provided as a reference tool only. Depending upon the type of tissues /tissue processing and reagents employed, the end user will need to optimize the final conditions for achieving an expected staining.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NB100-65125

NBP1-96778	Mouse IgG2a Isotype Control (M2A)
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NBL1-10816	FoxP1 Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-65125

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

