Product Datasheet

NRSF Antibody - BSA Free NB100-756

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-756

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-756



NB100-756


NRSF Antibody - BSA Free

Product Information Unit Size 100 ul Concentration 1.0 mg/ml Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0) Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.	INRSF Antibody - BSA Free	
Concentration 1.0 mg/ml Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0) Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry, Immunoprecipitation 1:500-1:2000, Western Blot (Negative) Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Product Information	
Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0) Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Unit Size	100 ul
Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0) Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Concentration	1.0 mg/ml
Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0) Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Storage	Store at 4C. Do not freeze.
Isotype IgG Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0) Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Clonality	Polyclonal
Purity Immunogen affinity purified Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0) Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Preservative	0.09% Sodium Azide
Buffer Tris-Citrate/Phosphate (pH 7.0 - 8.0) Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Isotype	IgG
Product Description Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Purity	Immunogen affinity purified
Host Rabbit Gene ID 5978 Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)
Gene Symbol REST Species Human Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Product Description	
Gene Symbol REST Species Human The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Host	Rabbit
Species	Gene ID	5978
Immunogen The immunogen recognized by this antibody maps to a region between residues 800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Gene Symbol	REST
800 and 850 of human RE1-silencing transcription factor using the numbering given in entry NP_005603.2 (GeneID 5978). Product Application Details Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Species	Human
Applications Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Immunogen	800 and 850 of human RE1-silencing transcription factor using the numbering
Western Blot (Negative) Recommended Dilutions Immunohistochemistry, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Product Application Details	
Immunohistochemistry-Paraffin 1:500-1:2000, Western Blot (Negative) Application Notes Western Blot Not recommended. Use rabbit anti-REST antibody NB100-757. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Applications	
Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue	Recommended Dilutions	
	Application Notes	Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue




Images

Immunohistochemistry-Paraffin: NRSF Antibody [NB100-756] - Sample: FFPE section of human ovarian carcinoma. Antibody: Affinity purified rabbit anti- REST used at a dilution of 1:1,000 (1ug/ml). Detection: DAB



Immunoprecipitation: NRSF Antibody [NB100-756] - Detection of human REST by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-REST antibody NB100-756 used for IP at 6 ug per reaction. For blotting immunoprecipitated REST, NB100-756 was used at 1 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



Publications

Kashiwagi K, Ishii J, Sakaeda M et al. Differences of molecular expression mechanisms among neural cell adhesion molecule 1, synaptophysin, and chromogranin A in lung cancer cells. Pathol Int 2012-04-01 [PMID: 22449227]



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB100-756

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-42048PEP NRSF Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-756

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

