

Product Datasheet

FEM1C Antibody NB100-784

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 7


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-784

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-784



NB100-784**FEM1C Antibody**

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA


Product Description	
Host	Goat
Gene ID	56929
Gene Symbol	FEM1C
Species	Mouse, Rat
Immunogen	Peptide with sequence C-HIPEKLETFVSLHR corresponding to C-Terminus according to NP_064562.1.

Product Application Details	
Applications	Western Blot, Peptide ELISA
Recommended Dilutions	Western Blot 0.3 - 1 ug/ml, Peptide ELISA Detection limit 1:8000
Application Notes	WB: Approx. 75 kDa band observed in heart and kidney lysates from mouse and rat (calculated MW of 68.6 kDa band according to mouse NP_775599.1).



Images

Western Blot: FEM1C Antibody [NB100-784] - Analysis of FEM1C in Hela lysate using NB100-784 at 2 ug/ ml.



Western Blot: FEM1C Antibody [NB100-784] - (0.5ug/ml) staining of rat kidney lysate (RIPA buffer, 35ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



Publications

Pritchard KA, Jing X, Teng M et al. Role of endoplasmic reticulum stress in impaired neonatal lung growth and bronchopulmonary dysplasia PloS one 2022-08-26 [PMID: 36018859]

Clift D, McEwan WA, Labzin LI et al. A Method for the Acute and Rapid Degradation of Endogenous Proteins. Cell. 2017-11-13 [PMID: 29153837]

de la Pisa IG, Cebrian C, Ortega JE et al. Cytokine pathway disruption in a mouse model of schizophrenia induced by Munc18-1a overexpression in the brain. J Neuroinflammation 2014-07-29 [PMID: 25069615]

Details:

CD11b antibody used for WB application at 1:1000 dilution in mouse model of schizophrenia induced by Munc18-1a overexpression/OE in the brain (see full text for sample preparation details). Figures 3a and 4a shows the WB of microglia /macrophage marker CD11bas single band of 160 kDa in cerebral cortex and striatum tissues of Munc18-OE and wild-type mice.

Koen YM, Sarma D, Hajovsky H et al. Protein Targets of Thioacetamide Metabolites in Rat Hepatocytes Chem Res Toxicol 2013-03-20 [PMID: 23465048]

Roehlecke C, Schumann U, Ader M et al. Stress Reaction in Outer Segments of Photoreceptors after Blue Light Irradiation. PLoS One. 2013-09-11 [PMID: 24039718]

Teng RJ, Jing X, Michalkiewicz T et al. Attenuation of endoplasmic reticulum stress by caffeine ameliorates hyperoxia-induced lung injury. Am. J. Physiol. Lung Cell Mol. Physiol. 2017-05-01 [PMID: 28213471]

Chan SL, Tan KO, Zhang L et al. F1Aalpha, a death receptor-binding protein homologous to the Caenorhabditis elegans sex-determining protein, FEM-1, is a caspase substrate that mediates apoptosis. J Biol Chem 1999-11-05 [PMID: 10542291]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-784

NB800-PC1	HeLa Whole Cell Lysate
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-784

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



