

Product Datasheet

RelA/NFkB p65 Antibody NB100-78423

Unit Size: 0.2 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-78423

Updated 12/20/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-78423



NB100-78423

RelA/NFkB p65 Antibody

Product Information	
Unit Size	0.2 ml
Concentration	0.5 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Protein A or G purified
Buffer	PBS (pH 7.2) and 0.2% Gelatin
Target Molecular Weight	65 kDa

Product Description	
Host	Rabbit
Gene ID	5970
Gene Symbol	RELA
Species	Human, Mouse, Rat
Specificity/Sensitivity	NF-kB p65
Immunogen	peptide-KLH, C-terminal

Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence 1:100-1:400, Immunoprecipitation 1:10-1:500, Immunohistochemistry-Paraffin 1:10-1:500
Application Notes	Each lot of this antibody is quality control tested by Western blotting. Western blotting, suggested working dilution(s): Use 10 ul per 5 ml antibody dilution buffer for each mini-gel. For immunofluorescence microscopy: Use a dilution range of 1:100~1:400. It is recommended that the reagent be titrated for optimal performance for each application.

Publications

de la Pisa IG, Cebrian C, Ortega JE et al. Cytokine pathway disruption in a mouse model of schizophrenia induced by Munc18-1a overexpression in the brain. *J Neuroinflammation* 2014-07-29 [PMID: 25069615] (WB, Mouse)

Details:

CD11b antibody used for WB application at 1:1000 dilution in mouse model of schizophrenia induced by Munc18-1a overexpression/OE in the brain (see full text for sample preparation details). Figures 3a and 4a shows the WB of microglia /macrophage marker CD11bas single band of 160 kDa in cerebral cortex and striatum tissues of Munc18-OE and wild-type mice.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-78423

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

