# **Product Datasheet**

# **EPB41L5 Antibody NB100-93392**

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-93392

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-93392



# NB100-93392


EPB41L5 Antibody

Product Information Unit Size	
Concentration 0.5 mg/ml Storage Store at -20C. Avoid freeze-thaw cycles.  Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA  Product Description Host Goat Gene ID 57669 Gene Symbol EPB41L5	
Storage Store at -20C. Avoid freeze-thaw cycles.  Clonality Polyclonal  Preservative 0.02% Sodium Azide  Isotype IgG  Purity Immunogen affinity purified  Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA  Product Description  Host Goat  Gene ID 57669  Gene Symbol EPB41L5	
Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA  Product Description Host Goat Gene ID 57669 Gene Symbol EPB41L5	
Preservative 0.02% Sodium Azide  Isotype IgG Purity Immunogen affinity purified  Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA  Product Description Host Goat Gene ID 57669 Gene Symbol EPB41L5	
Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA  Product Description Host Goat Gene ID 57669 Gene Symbol EPB41L5	
Purity Immunogen affinity purified  Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA  Product Description  Host Goat  Gene ID 57669  Gene Symbol EPB41L5	
Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA  Product Description  Host Goat  Gene ID 57669  Gene Symbol EPB41L5	
Product Description           Host         Goat           Gene ID         57669           Gene Symbol         EPB41L5	
Host         Goat           Gene ID         57669           Gene Symbol         EPB41L5	
Gene ID         57669           Gene Symbol         EPB41L5	
Gene Symbol EPB41L5	
-	
On a stars	
Species Human	
Immunogen Peptide with sequence C-ENLPQSPGTDQHD, from the internal region of the protein sequence according to NP_065960.2; NP_001171866.1; NP_001171868.1; NP_001317239.1.	
Product Application Details	
Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Peptide ELISA	
Recommended Dilutions  Western Blot 0.3-1 ug/ml, Flow Cytometry 10 ug/ml, Immunocytochemistry/ Immunofluorescence 10 ug/ml, Peptide ELISA Detection limit 1:8000	
Application Notes  WB: Approx. 80kDa band observed in lysates of cell line A431 and approx.  75kDa in lysates of cell line MCF7 (calculated MW of 81.9kDa according to NP_065960.2). Primary incubation 1 hour at room temperature.	




#### **Images**


Immunocytochemistry/Immunofluorescence: EPB41L5 Antibody [NB100-93392] - Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear, plasma membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).




Flow Cytometry: EPB41L5 Antibody [NB100-93392] - Flow cytometric analysis of paraformaldehyde fixed A549 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



Western Blot: EPB41L5 Antibody [NB100-93392] - (1ug/ml) staining of A431 (A) and (0.3ug/ml) MCF7 (B) cell lysate (35ug protein in RIPA buffer). Detected by chemiluminescence.



Immunocytochemistry/Immunofluorescence: EPB41L5 Antibody [NB100-93392] - Immunofluorescence analysis of paraformaldehyde fixed A549 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

# **Products Related to NB100-93392**

HAF017 Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

NBP2-30920PEP EPB41L5 Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-93392

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



