

Product Datasheet

TSLP Antibody - BSA Free NB110-55234

Unit Size: 0.05 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB110-55234

Updated 10/23/2024 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB110-55234



NB110-55234

TSLP Antibody - BSA Free

Product Information

Unit Size	0.05 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Peptide affinity purified
Buffer	PBS
Target Molecular Weight	18 kDa

Product Description

Host	Rabbit
Gene ID	85480
Gene Symbol	TSLP
Species	Human
Specificity/Sensitivity	Human TSLP has 2 isoforms, including isoform 1 (159aa, 18kD) and isoform 2 (63aa, 7kD). Mouse TSLP has one isoform (140aa, 16kD) and Rat TSLP also has one isoform (135aa, 16kD). NB110-55234 can detect only longer human isoform, but cannot detect shorter human isoform, mouse and rat.
Immunogen	Antibody was raised against a peptide corresponding to 19 amino acids near the center of human TSLP. The immunogen is located within the amino acids 40-90 of TSLP. Amino Acid Sequence: SGTKSTEFNNTVSCSNRPH

Product Application Details


Applications	Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence
Recommended Dilutions	Western Blot 0.25-4 ug/ml, ELISA 1:100-1:2000, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence 20 ug/mL, Immunohistochemistry-Paraffin 2.5 ug/ml, Immunofluorescence 20 ug/mL

Images

Western Blot: TSLP Antibody [NB110-55234] - EGF attenuates protein expression of PAR-2 and TSLP in DNCB-induced AD-like skin lesions in NC/Nga mice. Lysates were immunoblotted with anti-phospho-ERK, anti-ERK, anti-PAR-2, anti-TSLP, and anti-beta-actin antibodies. Cropped images from the different parts were displayed. Image collected and cropped by CiteAb from the following publication (www.nature.com/articles/s41598-018-30404-x) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: TSLP Antibody [NB110-55234] - Human Brain cells.




Immunohistochemistry: TSLP Antibody [NB110-55234] - EGF attenuates protein expression of PAR-2 and TSLP in DNCB-induced AD-like skin lesions in NC/ Nga mice. Paraffin sections of skin biopsies from DNCB-induced AD-like skin lesions were immunostained with anti-PAR-2 and anti-TSLP antibodies. Dark brown dots (Te red arrows) indicate positively stained cells. Images are representatives of groups. Scale bar, 100 μ m. Image collected and cropped by CiteAb from the following publication (www.nature.com/articles/s41598-018-30404-x) licensed under a CC-BY license.




Western Blot: TSLP Antibody [NB110-55234] - Total protein levels of TSLP in patients with AERD and subjects with CRS measured by using Western blotting of 3 representative samples per group. This image was submitted via customer Review.




Western Blot: TSLP Antibody [NB110-55234] - Western Blot Validation in Human Heart (Lane 1) and Human Prostate (Lane 2). Loading: 15 μ g of lysates per lane. Antibodies: TSLP NB110-55234 (4 μ g/mL), 1h incubation at RT in 5% NFDM/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.



Western Blot: TSLP Antibody [NB110-55234] - Western Blot Validation of TSLP with Human Recombinant protein. Loading: 30ng of TSLP partial human recombinant protein per lane. Antibodies: TSLP NB110-55234 (Lane 1: 0.25 ug/mL, Lane 2: 0.5 ug/mL, Lane 3: 1 ug/mL), 1h incubation at RT in 5% NFDM/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.



Immunohistochemistry-Paraffin: TSLP Antibody [NB110-55234] - Human brain tissue with TSLP antibody at 2.5 ug/mL.



Publications

Kanninen T, Tao L, Romero R et al. Thymic stromal lymphopoietin participates in the host response to intra-amniotic inflammation leading to preterm labor and birth Human immunology 2023-07-06 [PMID: 37422429] (WB, Human)

Chen C, Dai CY, Han F Et al. Interactions of thymic stromal lymphopoietin with interleukin-4 in adaptive immunity during Aspergillus fumigatus keratitis International journal of ophthalmology 2021-10-18 [PMID: 34667722] (WB, Mouse)

Kim YJ, Choi MJ, Bak DH et al. Topical administration of EGF suppresses immune response and protects skin barrier in DNCB-induced atopic dermatitis in NC/Nga mice Sci Rep 2018-08-09 [PMID: 30093649] (WB, Mouse)

Han Sang-Chul, Kang Gyeong-Jin, Ko Yeong-Jong et al. External Application of Fermented Olive Flounder (Paralichthys olivaceus) Oil Alleviates Inflammatory Responses in 2,4-Dinitrochlorobenzeneinduced Atopic Dermatitis Mouse Model. Toxicological Research 2012-01-01 [PMID: 24278605] (IF/IHC, WB, Mouse)

Han Sang-Chul, Kang Gyeong-Jin, Ko Yeong-Jong et al. Fermented fish oil suppresses T helper 1/2 cell response in a mouse model of atopic dermatitis via geneRation of CD4+CD25+Foxp3+ T cells. BMC Immunology 2012-01-01 [PMID: 22873180] (IF/IHC, WB, Mouse)

Buchheit KM, Cahill KN, Katz HR et al. Thymic stromal lymphopoietin controls prostaglandin D2 generation in patients with aspirin-exacerbated respiratory disease. J. Allergy Clin. Immunol. 2015-12-12 [PMID: 26691435]

Details:

TSLP antibody was used for WB analysis of lysates from nasal polyp samples of patients with aspirin-exacerbated respiratory disease/AERD (Fig 4; Fig E3). The antibody was tested for specificity using its corresponding blocking peptide NB110-55234PEP and was also compared with anti-TSLP antibody (ABT330 from Millipore; Fig E2).

Szicsz E, Molnar K, Lippai R et al. Peroxisome proliferator-activated receptor-γ and thymic stromal lymphopoietin are involved in the pathophysiology of childhood coeliac disease. Virchows Arch. 2014-09-04 [PMID: 25187315] (IHC-P, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB110-55234

NB110-55234PEP	TSLP Antibody Blocking Peptide
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-55234

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

