

# Product Datasheet

## EGFR [p Tyr1068] Antibody (EP38Y) NB110-56944

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB110-56944](http://www.novusbio.com/NB110-56944)

Updated 2/13/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB110-56944](http://www.novusbio.com/reviews/destination/NB110-56944)



**NB110-56944**

EGFR [p Tyr1068] Antibody (EP38Y)

<b>Product Information</b>	
<b>Unit Size</b>	0.1 ml
<b>Concentration</b>	This product is unpurified. The exact concentration of antibody is not quantifiable.
<b>Storage</b>	Store at -20C. Avoid freeze-thaw cycles.
<b>Clonality</b>	Monoclonal
<b>Clone</b>	EP38Y
<b>Preservative</b>	0.01% Sodium Azide
<b>Isotype</b>	IgG
<b>Purity</b>	Tissue culture supernatant
<b>Buffer</b>	49% PBS, 0.05% BSA and 50% Glycerol
<b>Target Molecular Weight</b>	175 kDa
<b>Product Description</b>	
<b>Host</b>	Rabbit
<b>Gene ID</b>	1956
<b>Gene Symbol</b>	EGFR
<b>Species</b>	Human, Mouse, Rat
<b>Reactivity Notes</b>	Reacts with: Mouse, Rat, Human
<b>Specificity/Sensitivity</b>	The antibody detects both EGFR phosphorylated on Tyrosine 1068 of the mature human isoform 1 (corresponding to Y1092 from the precursor form P00533-1/p170), and also unphosphorylated EGFR.
<b>Immunogen</b>	A synthetic phospho-peptide corresponding to residues surrounding Tyr1068 of human EGF receptor was used as immunogen.
<b>Notes</b>	Produced using Abcam's RabMab® technology. RabMab® technology is covered by the following U.S. Patents, No. 5,675,063 and/or 7,429,487.
<b>Product Application Details</b>	
<b>Applications</b>	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
<b>Recommended Dilutions</b>	Western Blot 1:1000-10000, Simple Western, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence 1:250-500, Immunoprecipitation 1:50, Immunohistochemistry-Paraffin 1:100-250
<b>Application Notes</b>	This product is useful for: Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry, Immunoprecipitation. In Western blot this antibody detects a band at approximately 175kDa.



**Immunohistochemistry Protocol for EGFR Antibody (NB110-56944)**

## Immunohistochemistry Protocol for Paraffin-embedded Tissues

**1. Solutions and reagents**

## 1.1. Xylene

1.2. Ethanol, anhydrous denatured, histological grade (100%, 95%, 70%)

## 1.3. Washing buffer:

TBST washing buffer: 1XTBS/0.1% Tween-20

To prepare stock solution of 10X TBS: add 24.2 g Trizma base and 80 g sodium chloride to 1L of dH<sub>2</sub>O. Adjust pH to 7.6.Working solution. 1XTBST/0.1% Tween-20: add 100ml 10XTBS to 900 ml dH<sub>2</sub>O. Add 1 ml Tween-20 and mix well.1.4. Distilled water (dH<sub>2</sub>O)

## 1.5. Antigen Retrieval Solution:

0.01M Sodium Citrate Buffer, pH 6.0

To prepare stock solutions:

Solution A. 0.1 M citric acid solution: dissolve 21.0 g of citric acid, monohydrate (C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>.H<sub>2</sub>O) in 1 liter of dH<sub>2</sub>O.Solution B. 0.1M sodium citrate solution: dissolve 29.4 g trisodium citrate dihydrate (C<sub>6</sub>H<sub>5</sub>Na<sub>3</sub>O<sub>7</sub>.2H<sub>2</sub>O) in 1 liter of dH<sub>2</sub>O.Working solution: Add 9 ml of Stock solution A and 41 ml of stock solution B to 450 ml of dH<sub>2</sub>O. Adjust pH to 6.0.

## 1.6. 3% Hydrogene Peroxide

## 1.7. Blocking buffer:

PBS (Dulbeccos Phosphate Buffered Salts, 1X, catalog #21-031-CV from Mediatech, Inc.) + 10% serum (serum origin depends on the host of the secondary antibody)

## 1.8. Hematoxylin QS (catalog #H-3404 from Vector Laboratories, Inc.)

## 1.9. Permanent Mounting medium (VectaMount, catalog# H-5000 Vector Laboratories, Inc.)

**2. Protocol****2.1. Deparaffinization/Rehydration**

2.1.1. Heat slides in an oven at 65C for 1 hour.

2.1.2. De-paraffinize/hydrate using the following series of washes: two Xylene washes (5 min each), followed by two 100% ethanol rinses (5 min each), followed by 95% ethanol, 70% ethanol, 50% ethanol, 30% ethanol, followed by H<sub>2</sub>O and a TBST wash for 5 min on a shaker.**2.2. Antigen Retrieval**

2.2.1. Immerse slides into staining dish containing Antigen Retrieval Solution.

2.2.2. Place covered staining dish into the rice cooker. Add 120 ml of dH<sub>2</sub>O.

2.2.3. When cook is turned to warm (about 20 to 30 min), unplug the cooker and remove the staining dish to the bench top.

2.2.4. Allow to cool down, without cover, for 20 min.

**2.3. Staining**

2.3.1. Wash slides with TBST for 5 min on a shaker.

2.3.2. Inactivate endogenous peroxidase by covering tissue with 3% hydrogen peroxide for 10 min.

2.3.3. Wash slides three times with TBST (3 min each on a shaker).

2.3.4. Block slides with the blocking solution for 1 hour.

2.3.5. Dilute primary antibody in the blocking buffer per recommendation on the data sheet.

2.3.6. Apply primary antibody to each section and incubate overnight in the humidified chamber (4C).

2.3.7. Wash slides three times with TBST (3 min each on a shaker).

2.3.8. Apply to each section secondary HRP-conjugated anti-rabbit antibody diluted in the blocking solution per manufacturers recommendation; incubate for 1 hour at room temperature.

2.3.9. Wash slides three times with TBST (3 min each on a shaker).

2.3.10. Add freshly prepared DAB substrate to the sections.

2.3.11. Incubate tissue sections with the substrate at room temperature until suitable staining develops (generally 2 to 5 min).

2.3.12. Rinse sections with water.

2.3.13. Counterstain with Hematoxylin.

2.3.14. Rinse sections with water.

2.3.15. Dehydrate samples using two rinses with 100% Ethanol (20 dips per rinse) followed by two rinses with Xylene (30 dips per rinse).

2.3.16. Mount coverslips on slides using Permount medium.



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB110-56944](http://www.novusbio.com/reviews/submit/NB110-56944)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

