

# Product Datasheet

## Enolase 2/Neuron-specific Enolase Antibody NB110-58870

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 2 Publications: 5**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB110-58870](http://www.novusbio.com/NB110-58870)

Updated 4/13/2025 v.20.1

**Earn rewards for product  
reviews and publications.**

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB110-58870](http://www.novusbio.com/reviews/destination/NB110-58870)



**NB110-58870****Enolase 2/Neuron-specific Enolase Antibody****Product Information**

<b>Unit Size</b>	0.1 ml
<b>Concentration</b>	This product is unpurified. The exact concentration of antibody is not quantifiable.
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Polyclonal
<b>Preservative</b>	0.035% Sodium Azide
<b>Isotype</b>	IgG
<b>Purity</b>	Unpurified
<b>Buffer</b>	Supplied as serum
<b>Target Molecular Weight</b>	47 kDa

**Product Description**


<b>Host</b>	Rabbit
<b>Gene ID</b>	2026
<b>Gene Symbol</b>	ENO2
<b>Species</b>	Human, Mouse, Rat
<b>Marker</b>	Neuronal Marker
<b>Immunogen</b>	Recombinant full length human NSE purified from E. coli. [UniProt# P09104]

**Product Application Details**


<b>Applications</b>	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Knockdown Validated
<b>Recommended Dilutions</b>	Western Blot 1:2000, Simple Western 1:10000, Immunohistochemistry 1:500, Immunocytochemistry/ Immunofluorescence 1:500, Knockdown Validated
<b>Application Notes</b>	<p>This NSE antibody is useful for Immunocytochemistry/Immunofluorescence and Western Blot.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See <a href="#">Simple Western Antibody Database</a> for Simple Western validation: Tested in Human Brain lysate 0.05 mg/mL, separated by Size, antibody dilution of 1:10,000. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>

**Images**


Western Blot: Enolase 2/Neuron-specific Enolase Antibody [NB110-58870] - Analysis of different tissue and cell lysates using rabbit pAb to neuron specific enolase (NSE), NB110-58870, dilution 1:5,000 in green: [1] protein standard (red), [2] rat brain, [3] rat spinal cord, [4] mouse brain, [5] mouse spinal cord, [6] NIH-3T3, [7] HEK293, [8] HeLa, [9] SH-SY5Y, and [10] C6 cells. A single band at about 47kDa corresponds to the NSE protein, seen only in extracts containing neurons or neuronal lineage cells.




**Immunocytochemistry/Immunofluorescence: Enolase 2/Neuron-specific Enolase Antibody [NB110-58870]** - Analysis of mixed cortical neuron-glia cell culture from E20 rat stained with rabbit pAb to neuron specific enolase (NSE), NB110-58870, dilution 1:500 in red, and costained with chicken pAb to GFAP, dilution 1:5,000 in green. The blue is Hoechst staining of nuclear DNA. the NSE antibody labels protein expressed in neuronal cells, while the GFAP antibody stains intermediate filaments in astrocytic and certain other glial cells.




**Western Blot: Enolase 2/Neuron-specific Enolase Antibody [NB110-58870]** - Blot of rat spinal cord probed with rabbit antibody to NSE. The antibody stains a single sharp band corresponding to NSE at about 47kDa.




**Western Blot: Enolase 2/Neuron-specific Enolase Antibody [NB110-58870]** - Western blot analysis of extracts from mouse brain using NSE antibody (NB110-58870, 1:100).




**Immunocytochemistry/Immunofluorescence: Enolase 2/Neuron-specific Enolase Antibody [NB110-58870]** - Shows the human embryonic kidney cells line 293, which express many neuronal proteins (1). The red channel shows staining which recognizes all of these 293 cells. The green channels shows staining for another neuronal marker to UCHL1 using NB110-58869. This neuronal gene is apparently activated in a cell density dependent fashion and at this stage only a few cells express this protein. However all cells that express NSE also express UCHL1.



Immunocytochemistry/Immunofluorescence: Enolase 2/Neuron-specific Enolase Antibody [NB110-58870] - Confocal immunofluorescent analysis of 3T3 cells using NSE antibody (NB110-58870, 1:5). An Alexa Fluor 488-conjugated Goat to rabbit IgG was used as secondary antibody (green, A). Actin filaments were labeled with Alexa Fluor 568 phalloidin (red, B). DAPI was used to stain the cell nuclei (blue, C).



Simple Western: Enolase 2/Neuron-specific Enolase Antibody [NB110-58870] - Simple Western lane view shows a specific band for Enolase 2/Neuron specific enolase in 0.05 mg/ml of Human Brain lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



## Publications

Powers RM, Daza R, Koehler AE et al. Growth cone macropinocytosis of neurotrophin receptor and neuritogenesis are regulated by neuron navigator 1 Molecular biology of the cell 2022-06-01 [PMID: 35352947] (ICC/IF, Human)

Details:

Dilution used for ICC 1:250

Niture S, Tricoli L, Qi Q Et al. MicroRNA miR-99b-5p Targets mTOR/AR axis, Induces Autophagy, and Inhibits Prostate Cancer Cell Proliferation Tumour Biol 2022-07-11 [PMID: 35811549]

Gong W, Zhang S, Zong Y et al. hiPS Cell-Derived Neurons for High-Throughput Screening Methods Mol. Biol. 2019-05-25 [PMID: 31124122] (ICC/IF, Human)

Modarai S, Gupta A, Opdenaker L et al. The anti-cancer effect of retinoic acid signaling in CRC occurs via decreased growth of ALDH+ colon cancer stem cells and increased differentiation of stem cells Otolaryngol Head Neck Surg 2018-10-16 [PMID: 30410666] (WB, Human)

Jung Y, Cackowski FC, Yumoto K, Decker A. CXCL12gamma Promotes Development of Metastatic Castration Resistant Prostate Cancer by Induction of Cancer Stem Cell and Neuroendocrine Phenotypes. Cancer Res. 2018-02-05 [PMID: 29431639] (WB, Human)



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB110-58870**

---

NBL1-10268	Enolase 2/Neuron-specific Enolase Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB110-58870](http://www.novusbio.com/reviews/submit/NB110-58870)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

