

Product Datasheet

Granzyme B Antibody NB110-59915

Unit Size: 6 ml

Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB110-59915

Updated 7/19/2023 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB110-59915



NB110-59915**Granzyme B Antibody**

Product Information	
Unit Size	6 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.4), 0.2% BSA, Tween-20


Product Description	
Description	<p>This antibody is useful in the localization of granzyme B containing lytic granules and for the characterization of activated CTLs or NK cells.</p> <p>This product has a prediluted size available. If a 6mL or 7mL pre-diluted size is purchased this product can be used Neat (without further dilution). Use recommended dilution for the concentrated vials only.</p>
Host	Rabbit
Gene ID	3002
Gene Symbol	GZMB
Species	Human
Specificity/Sensitivity	This antibody reacts with a 32 kD protein.
Immunogen	A synthetic peptide from the N-terminus of human Granzyme B.

Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry 1:10-1:500, Immunohistochemistry-Paraffin
Application Notes	IHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT.




Images

Immunohistochemistry-Paraffin: Granzyme B Antibody [NB110-59915] - Formalin fixed paraffin embedded human Hodgkin's lymphoma stained with Granzyme B antibody (NB110-59915).



Immunohistochemistry-Paraffin: Granzyme B Antibody [NB110-59915] - ab4069 at 1/20 dilution staining rat spinal cord tissue sections by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). The tissue was formaldehyde fixed and blocked with serum prior to incubation with the antibody for 2 hours. A biotinylated goat anti-rabbit IgG was used as the secondary.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-59915

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

