

Product Datasheet

LMO2 Antibody (1A9-3B11) - BSA Free NB110-78626

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB110-78626

Updated 4/22/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB110-78626




NB110-78626

LMO2 Antibody (1A9-3B11) - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	1A9-3B11
Preservative	0.02% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	24 kDa
Product Description	
Host	Mouse
Gene ID	4005
Gene Symbol	LMO2
Species	Human, Mouse
Reactivity Notes	Use in Mouse reported in scientific literature (PMID:34382935).
Immunogen	Recombinant human LMO2. [UniProt# P25791]
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP), ICC/IF (Negative)
Recommended Dilutions	Western Blot 0.5 ug/ml, Flow Cytometry 1-2 ug/ml per million cells, Immunohistochemistry 1:100-1:500, Immunoprecipitation 1:10-1:500, Immunohistochemistry-Paraffin 1:100-1:500, ICC/IF (Negative), Chromatin Immunoprecipitation (ChIP) 1:10-1:500
Application Notes	This LMO2 antibody is useful for Western blot, Immunoprecipitation, Chromatin Immunoprecipitation and Immunohistochemistry on paraffin-embedded sections. In WB a band can be seen at ~24 kDa. The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.

Images


Western Blot: LMO2 Antibody (1A9-3B11) [NB110-78626] - Detection of LMO2 in Ramos cell lysate.




Immunohistochemistry-Paraffin: LMO2 Antibody (1A9-3B11) [NB110-78626] - IHC analysis of formalin fixed paraffin-embedded (FFPE) human tonsil using LMO2 antibody at 1:500 on a Bond Rx autostainer (Leica Biosystems). The assay involved 20 minutes of heat induced antigen retrieval (HIER) using 10mM sodium citrate buffer (pH 6.0) and endogenous peroxidase quenching with peroxide block. The sections were incubated with primary antibody for 30 minutes and Bond Polymer Refine Detection (Leica Biosystems) with DAB was used for signal development followed by counterstaining with hematoxylin. Whole slide scanning and capturing of representative images was performed using Aperio AT2 (Leica Biosystems). Nuclear with some cytoplasmic staining was observed. Staining was performed by Histowiz.




Flow Cytometry: LMO2 Antibody (1A9-3B11) [NB110-78626] - An intracellular stain was performed on Ramos cells with LMO2 Antibody (1A9-3B11) NB110-78626 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1.0 ug/mL for 30 minutes at room temperature, followed by Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Dylight 550 (35503, Thermo Fisher).



Flow Cytometry: LMO2 Antibody (1A9-3B11) [NB110-78626] - An intracellular stain was performed on THP-1 cells with LMO2 Antibody (1A9-3B11) NB110-78626 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1.0 ug/mL for 30 minutes at room temperature, followed by Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Dylight 550 (35503, Thermo Fisher).



Immunohistochemistry: LMO2 Antibody (1A9-3B11) - BSA Free [NB110-78626] - Immunohistochemistry labeling of LMO2 (green) in tonsil tissue.



Publications

Seo S, Hong N, Song J et al. Polymer Thin Film Promotes Tumor Spheroid Formation via JAK2-STAT3 Signaling Primed by Fibronectin-Integrin $\alpha 5$ and Sustained by LMO2-LDB1 Complex Biomedicines 2022-10-24 [PMID: 36359204] (WB, Human)

Wang W, Meng Y, Chen Y et al. A comprehensive analysis of LMO2 pathogenic regulatory profile during T-lineage development and leukemic transformation Oncogene 2022-07-18 [PMID: 35851847] (Chemotaxis, Mouse)

Hirano KI, Hosokawa H, Koizumi M Et al. LMO2 is essential to maintain the ability of progenitors to differentiate into T-cell lineage in mice eLife 2021-08-12 [PMID: 34382935] (Chemotaxis, WB, Mouse)

Natkunam, Y et al. The oncoprotein LMO2 is expressed in normal germinal-center B cells and in human B-cell lymphomas. Blood;109(4):1636-42. 2007-02-15 [PMID: 17038524] (IHC-P, WB, Human)

Procedures

Western Blot protocol for LMO2 Antibody (NB110-78626)

LMO2 Antibody (1A9-3B11):

Western Blot Protocol

1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 40 ug of total protein per lane.
2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
3. Rinse membrane with dH₂O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
4. Rinse the blot in TBS for approximately 5 minutes.
5. Block the membrane using 5% NFD_M + 1% BSA in TBS + Tween, 1 hour at RT.
6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
7. Dilute the rabbit anti-LMO2 primary antibody (NB 110-78626) in blocking buffer and incubate 1 hour at room temperature.
8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
9. Apply the diluted mouse-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturers instructions (Pierce ECL).

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB110-78626

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NB110-78626C	LMO2 Antibody (1A9-3B11) [DyLight 650]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-78626

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

