Product Datasheet

Survivin [p Ser20] Antibody - BSA Free NB110-92717SS

Unit Size: 0.025 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB110-92717

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB110-92717



NB110-92717SS

Survivin [p Ser20] Antibody - BSA Free

Product Information	
0.025 ml	
1.25 mg/ml	
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Polyclonal	
0.1% Sodium Azide	
IgG	
Immunogen affinity purified	
PBS, 30% Glycerol	
16 kDa	
Product Description	
Rabbit	
332	
BIRC5	
Human, Mouse, Rat	
Mouse reactivity reported in scientific literature (PMID: 22814318). Rat reactivity reported in scientific literature (PMID: 17612487). Human reactivity reported in scientific literature (PMID: 24069188).	
This Survivin [p Ser20] Antibody was developed against a synthetic peptide surrounding the phosphorylated serine 20 of the human Survivin protein. [Swiss-Prot #O15392]	
Product Application Details	
Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry	
Western Blot reported in scientific literature (PMID 24069188), Immunohistochemistry 1:10-1:500. Use reported in scientific literature (PMID 22814318), Immunocytochemistry/ Immunofluorescence 1:100	
The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.	

www.novusbio.com




Images

Western Blot: Survivin [p Ser20] Antibody [NB110-92717] - Detection of Survivin [Ser20] in phosphorylated recombinant protein (+), but not in unphosphorylated protein (-) using [NB110-92717]. Band detected at slightly higher than the predicted molecular weight of 16 kDa.

in Hela cells with DyLight 488 (green). Nuclei and alpha-tubulin were

counterstained with DAPI (blue) and DyLight 550 (red).



Immunocytochemistry/Immunofluorescence: Survivin [p Ser20] Antibody [NB110-92717] - Survivin [p Ser20] Antibody (NB110-92717) was tested

Publications

Garlapati C, Joshi S, Bhattarai S et al. PLK1 and AURKB phosphorylate survivin differentially to affect proliferation in racially distinct triple-negative breast cancer Cell Death & Disease 2023-01-10 [PMID: 36627281] (Immunohistochemistry)

Benaiges E, Ceperuelo-MallafrE V, Madeira A et al. Survivin drives tumor-associated macrophage reprogramming: a novel mechanism with potential impact for obesity Cellular oncology (Dordrecht) 2021-03-12 [PMID: 33710603]

Knapp Charles F, Sayegh Zena, Schell Michael J et al. Expression of CXCR4, E-cadherin, Bcl-2, and survivin in Merkel cell carcinoma: an immunohistochemical study using a tissue microarray. The American Journal of Dermatopathology 2012-01-01 [PMID: 22814318] (IF/IHC, Mouse)

Muschol-Steinmetz C, Friemel A, Kreis NN et al. Function of Survivin in Trophoblastic Cells of the Placenta. PLoS One. 2013-09-19 [PMID: 24069188] (WB, Human)

Dohi, T et al. Compartmentalized phosphorylation of IAP by protein kinase A regulates cytoprotection. Mol Cell;27 (1):17-28. 2007-07-06 [PMID: 17612487] (WB, Rat)



Procedures

Immunocytochemistry/Immunofluorescence Protocol for Survivin [p Ser20] antibody (NB110-92717) Survivin [p Ser20] Antibody:

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.

2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.

3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.

4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.

5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.

6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.

7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.

8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,0000 and incubate for 10 minutes. Wash a third time for 10 minutes.

9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-92717

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

