# **Product Datasheet**

# PSTAIR Antibody (PSTAIR) NB120-10345

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB120-10345

Updated 10/23/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB120-10345




# NB120-10345

PSTAIR Antibody (PSTAIR)	
Product Information	
Unit Size	0.1 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSTAIR
Preservative	0.1% Sodium Azide
Isotype	IgG1
Purity	Unpurified
Buffer	Raw ascites
Product Description	
Host	Mouse
Species	Human, Mouse, Avian, Plant, Primate, Xenopus
Reactivity Notes	Cross-reacts with human, monkey, mouse, Japanese quail, fish (goldfish, carp, eel, starfish, and amago salmon), Xenopus, cliliates, and plants (lily and onion). Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.
Specificity/Sensitivity	PSTAIR (PSTAIR)
Immunogen	Synthetic peptide: EGVPSTAIREISLLKE conjugated to BSA.
Product Application Details	
Applications	Western Blot, ELISA, Immunoprecipitation
Recommended Dilutions	Western Blot 1:100-1:2000, ELISA 1:100-1:2000, Immunoprecipitation 1:10-1:500
Application Notes	ELISA: Use at an assay dependent dilution. IP: Use at an assay dependent dilution. It cannot precipitate p34cdc2; when it is complexed with cyclin B.WB: Use at a dilution of 1/4000 (determined by indirect blotting using COS-7 cell extract). The antibody recognises 31-34kD proteins (1-4) bands.Monoclonal Anti-PSTAIR may be used for the localization of cyclin-dependent kinases, containing the PSTAIR motif.Not tested in other applications.Optimal dilutions/concentrations should be determined by the end user.



## **Images**

Western Blot: PSTAIR Antibody (PSTAIR) [NB120-10345] - Cell line lysates were separated on SDS-PAGE and probed with 1:500 Monoclonal Anti-PSTAIR Clone:PSTAIR. The antibody was developed using Goat Anti-Mouse IgG-Peroxidase and a chemiluminescent substrate. Lanes 1. HEK-293T 2. HeLa 3. SK-N-SH 4. SH-SY5Y 5. U87 6. COS7 7. PC12 8. Neuro-2a 9. Arabidopsis





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com
Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NB120-10345**

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB120-10345

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

