

Product Datasheet

EWSR1 Antibody - BSA Free

NB200-183

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB200-183

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB200-183



NB200-183

EWSR1 Antibody - BSA Free

Product Information

Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)

Product Description


Host	Rabbit
Gene ID	2130
Gene Symbol	EWSR1
Species	Human
Immunogen	A synthetic peptide that maps to a region between residues 600 and the C-terminus (residue 656) of human Ewing sarcoma breakpoint region 1 using the numbering given in SwissProt entry Q01844 (GenelD 2130).

Product Application Details


Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot 1:10000 - 1:25000, Immunoprecipitation 2-10 ug/mg lysate

Images

Western Blot: EWSR1 Antibody [NB200-183] - Detection of Human and Mouse EWS by Western Blot. Samples: Whole cell lysate (15 and 50 ug) from HeLa and mouse NIH3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-EWS antibody NB200-183 used for WB at 0.04 ug/ml. Detection: Chemiluminescence with an exposure time of 3 seconds.



Immunoprecipitation: EWSR1 Antibody [NB200-183] - Detection of human EWSR1 by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-EWSR1 antibody NB200-183 used for IP at 6 ug per reaction. EWSR1 was also immunoprecipitated by rabbit anti-EWSR1 antibody NB200-182. For blotting immunoprecipitated EWSR1, NB200-183 was used at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 3 seconds.



Publications

Sohn EJ, Park J, Kang SI et al. Accumulation of pre-let-7g and downregulation of mature let-7g with the depletion of EWS. Biochem Biophys Res Commun 2012-09-01 [PMID: 22910415]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB200-183

NBL1-10371	EWSR1 Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB200-183

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

