

Product Datasheet

DDX6 Antibody NB200-191

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 1 **Publications: 43**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB200-191

Updated 11/16/2022 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB200-191



NB200-191

DDX6 Antibody


Product Information	
Unit Size	0.1 ml
Concentration	0.2 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	TBS, 0.1% BSA

Product Description	
Host	Rabbit
Gene ID	1656
Gene Symbol	DDX6
Species	Human, Mouse, Hamster
Reactivity Notes	Hamster reactivity reported from a verified customer reivew.
Immunogen	The immunogen recognized by NB200-191 maps to a region between residues 1 and 50 of human DEAD (Asp-Glu-Ala-Asp) box polypeptide 6 using the numbering given Swiss-Prot entry P26196 (GeneID 1656).


Product Application Details	
Applications	Western Blot, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:5000-1:15000, Immunohistochemistry 1:100-1:500, Immunocytochemistry/Immunofluorescence, Immunoprecipitation 2-5 ug/mg lysate, Immunohistochemistry-Paraffin 1:100-1:500
Application Notes	ICC/IF was reported in scientific literature (PMID: 25800057) and a verified customer review. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Epitope retrieval with Tris-EDTA pH 9.0 is recommended for FFPE tissue sections.

Images


Western Blot: DDX6 Antibody [NB200-191] - Western blot analysis of LBCs show increased Staufen (STAU1) (NBP1-33202) levels compared with normal controls. DDX6 (NB200-191) levels are unchanged. HD and SCA3 patient (polyQ expanded) FBs were used as additional controls. Four normal and five SCA2 FBs, and two normal and three SCA2 LBCs were used. Image collected and cropped by CiteAb from the following publication (<http://pubmed.ncbi.nlm.nih.gov/30194296/>) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: DDX6 Antibody [NB200-191] - In normal condition, cells were fixed and stained for DDX6, that usually punctuate in process bodies (PBs). Primary Antibody rabbit-anti-DDX6 diluted 1:1000. Image submitted by a verified customer review.



Immunohistochemistry-Paraffin: DDX6 Antibody [NB200-191] - Section of human colon carcinoma (left) and mouse renal cell carcinoma (right). Antibody: Affinity purified rabbit anti-DDX6 used at a dilution of 1:200 (1ug/ml). Detection: DAB.



Immunoprecipitation: DDX6 Antibody [NB200-191] - Detection of human DDX6 by Western Blot and Immunoprecipitation. Samples: Whole cell lysate (15 and 50 ug) from HeLa cells. Antibodies: Affinity purified rabbit anti-DDX6 antibody NB200-191 used at 0.04 ug/ml for WB (A and B) and at 3 ug/mg lysate for IP. DDX6 was also immunoprecipitated using rabbit anti-DDX6 antibody NB200-192, but not with 'DDX6 Ab', which is to another epitope from DDX6. Normal rabbit IgG was used as a negative control. Detection: Chemiluminescence with an exposure time of 1 second (A & B).



Publications

Venkataramanan S, Gadek M, Calviello L et al. DDX3X and DDX3Y are redundant in protein synthesis RNA 2021-12-01 [PMID: 34535544]

Heim A, Niedermeier ML, Stengel F, Mayer TU The translation regulator Zar1l controls timing of meiosis in Xenopus oocytes Development (Cambridge, England) 2022-11-01 [PMID: 36278895]

Mehravar M, Kumar Y, Olshansky M et al. MOV10 facilitates messenger RNA decay in an N6-methyladenosine (m6A) dependent manner to maintain the mouse embryonic stem cells state bioRxiv 2021-08-12

Sung CC, Poll B, Lin SH et al. Early Molecular Events Mediating Loss of Aquaporin-2 during Ureteral Obstruction in Rat Journal of the American Society of Nephrology : JASN 2022-08-02 [PMID: 35918145] (IHC-P, Rat)

Details:

Dilutions: 1:500

Hallacli E, Kayatekin C, Nazeen S et al. The Parkinson's disease protein alpha-synuclein is a modulator of processing bodies and mRNA stability Cell 2022-06-09 [PMID: 35688132] (ICC/IF, WB, Human)

Shah A, Bhandari R IP6K1 upregulates the formation of processing bodies by influencing protein-protein interactions on the mRNA cap Journal of cell science 2021-11-29 [PMID: 34841428]

Somasekharan SP, Gleave M SARS-CoV-2 nucleocapsid protein interacts with immunoregulators and stress granules and phase separates to form liquid droplets FEBS letters 2021-11-15 [PMID: 34780058] (ICC/IF, Human)

Rink MR, Baptista MAP, Flomm FJ Et al. Concatemeric Broccoli reduces mRNA stability and induces aggregates PloS one 2021-08-04 [PMID: 34347781] (ICC/IF)

Shah A, Bhandari R IP6K1 upregulates the formation of processing bodies by promoting proteome remodeling on the mRNA cap bioRxiv 2020-07-14 [PMID: 34841428] (WB, ICC/IF, Human)

Cougnoux A, Drummond RA, Fellmeth M, et al. GC content shapes mRNA storage and decay in human cells Elife 2019-12-19 [PMID: 31855182] (WB, ICC/IF, Human)

Sharma NR, Majerciak V, Kruhlak MJ et al. KSHV RNA-binding protein ORF57 inhibits P-body formation to promote viral multiplication by interaction with Ago2 and GW182 Nucleic Acids Res. 2019-08-10 [PMID: 31400113] (WB, Human)

Hubstenberger A, Courel M, Benard M, Souquere S et al. P-Body Purification Reveals the Condensation of Repressed mRNA Regulons Mol. Cell 2017-10-05 [PMID: 28965817] (ICC/IF, Human)

More publications at <http://www.novusbio.com/NB200-191>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB200-191

NBL1-09810	DDX6 Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB200-191

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

