

Product Datasheet

NCOA6 Antibody - BSA Free

NB200-335

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 8

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB200-335

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB200-335



NB200-335

NCOA6 Antibody - BSA Free

Product Information	
Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)


Product Description	
Host	Rabbit
Gene ID	23054
Gene Symbol	NCOA6
Species	Human
Immunogen	A synthetic peptide that maps to a region between residues 1050 and 1100 of human Activating Signal Cointegrator-2 using the numbering given in Swiss-Prot entry Q14686 (GeneID 23054).

Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:2000 - 1:10000, Immunohistochemistry 1:500 - 1:2000, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:500 - 1:2000
Application Notes	Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.




Images


Western Blot: NCOA6 Antibody [NB200-335] - Detection of Human ASC2 by Western Blot. Samples: Whole cell lysate (5, 15 and 50 ug) from HeLa cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-ASC2 antibody NB200-335 used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



Immunohistochemistry-Paraffin: NCOA6 Antibody [NB200-335] - FFPE section of human colon carcinoma. Antibody: Affinity purified rabbit anti-ASC2 used at a dilution of 1:1,000 (1ug/ml). Detection: DAB



Immunoprecipitation: NCOA6 Antibody [NB200-335] - Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-ASC2 antibody used for IP at 6 ug per reaction. ASC2 was also immunoprecipitated by a previous lot of this antibody and rabbit anti-ASC2 antibody. For blotting immunoprecipitated ASC2, was used at 1 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



Publications

Oh GS, Kim SR, Lee ES et al. Regulation of Hepatic Gluconeogenesis by Nuclear Receptor Coactivator 6 Molecules and Cells 2022-04-30 [PMID: 35258009] (Western Blot, Block/Neutralize)

Kopanja D, Pandey A, Huang S et al. p19Arf Inhibits Aggressive Progression of H-ras driven Hepatocellular Carcinoma. Carcinogenesis 2017-12-08 [PMID: 29228217] (IF/IHC, Human)

Chen WD, Fu X, Dong B et al. Neonatal activation of the nuclear receptor CAR results in epigenetic memory and permanent change of drug metabolism in mouse liver. Hepatology 2012-10-01 [PMID: 22488010]

Vicent GP, Nacht AS, Font-Mateu J et al. Four enzymes cooperate to displace histone H1 during the first minute of hormonal gene activation. Genes Dev 2011-04-01 [PMID: 21447625]

Lee S, Kim DH, Goo YH et al. Crucial roles for interactions between MLL3/4 and INI1 in nuclear receptor transactivation. Mol Endocrinol 2009-05-01 [PMID: 19221051]

Garapaty S, Xu CF, Trojer P et al. Identification and characterization of a novel nuclear protein complex involved in nuclear hormone receptor-mediated gene regulation. J Biol Chem 2009-03-01 [PMID: 19131338]

Lee J, Saha PK, Yang QH et al. Targeted inactivation of MLL3 histone H3-Lys-4 methyltransferase activity in the mouse reveals vital roles for MLL3 in adipogenesis. Proc Natl Acad Sci U S A 2008-12-01 [PMID: 19047629]

Cho YW, Hong T, Hong S et al. PTIP associates with MLL3- and MLL4-containing histone H3 lysine 4 methyltransferase complex. J Biol Chem 2007-07-01 [PMID: 17500065]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB200-335

NB800-PC9	HeLa Nuclear Cell Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB200-335

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

