

Product Datasheet

Caspase-8 Antibody (FLICE 4-1-20) NB500-208

Unit Size: 0.2 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB500-208

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB500-208



NB500-208**Caspase-8 Antibody (FLICE 4-1-20)**

Product Information	
Unit Size	0.2 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at 4C. Do not freeze.
Clonality	Monoclonal
Clone	FLICE 4-1-20
Preservative	0.1% Sodium Azide
Isotype	IgG1 Kappa
Purity	Unpurified
Buffer	Ascites
Product Description	
Host	Mouse
Gene ID	841
Gene Symbol	CASP8
Species	Human, Mouse
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 29069785).
Immunogen	Full-length recombinant human Caspase 8 [UniProt# Q14790]
Product Application Details	
Applications	Western Blot
Recommended Dilutions	Western Blot 1:1000
Application Notes	This Caspase 8 (FLICE 4-1-20) antibody is useful for Western Blot where a band is seen at 55 kDa representing the inactive form of Caspase 8 (see protocol for additional information). We also have reports that the antibody detects the active (pro) form of Caspase 8.




Images

Western Blot: Caspase-8 Antibody (FLICE 4-1-20) [NB500-208] - CaSki cells were mock-infected or exposed to each of the indicated viruses (MOI ~ 1 pfu/cell) or 10 uM staurosporine and 4 h pi, cellular lysates were prepared and analyzed by immunoblotting for Caspase-8. Image collected and cropped by CiteAb from the following publication (<https://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1006766>) licensed under a CC-BY license.



Western Blot: Caspase 8 Antibody (FLICE 4-1-20) [NB500-208] - Detection of Caspase 8 (59 kDa) from HEK293 cell extract using NB500-208 (1:500). Lanes 1 and 2 contain inactive and active Caspase 8, respectively.



Publications

Martin ACBM, Fuzer AM, Becceneri AB et al. [10]-gingerol induces apoptosis and inhibits metastatic dissemination of triple negative breast cancer in vivo. *Oncotarget*. 2017-09-22 [PMID: 29069785] (WB, Mouse)

Procedures

Protocol specific for Caspase 8 Antibody (NB500-208)

Caspase-8 Antibody (FLICE 4-1-20):

Western Protocol

- 1) Run a 10-*15% SDS polyacrylamide gel, loading ~20 ug of cell extract per lane.
- 2) Transfer the proteins to a membrane.
- 3) Block the membrane in PT-T20 [20 mM Tris-HCl, pH 7.4 / 150 mM NaCl/0.5% Tween 20] + 5% NFDM [non-fat dry milk] , for 3 hour at room temperature (RT), shaking gently.
- 4) Rinse the membrane twice with PT-T20.
- 5) Incubate the membrane in anti-Caspase 8 [NB 500-208], diluted 1:500-1:1,000 in PT-T20 + 5% NFDM, for 60 minutes at RT.
- 6) Wash the membrane for 15 minutes, 3 times, in PT-T20 at RT.
- 7) Incubate the membrane in anti-mouse conjugated to HRP (secondary antibody), diluted in PT-T20 + 5% NFDM for 60 minutes at RT.
- 8) Wash the membrane for 15 minutes, 3 times, in PT-T20 at RT.
- 9) Develop membrane in chemiluminescent reagents, as instructed by kit-vendor.

*Positive control(s): Human kidney 293 cells

Western Protocol

- 1) Run a 10-*15% SDS polyacrylamide gel, loading ~20 ug of cell extract per lane.
- 2) Transfer the proteins to a membrane.
- 3) Block the membrane in PT-T20 [20 mM Tris-HCl, pH 7.4 / 150 mM NaCl/0.5% Tween 20] + 5% NFDM [non-fat dry milk] , for 3 hour at room temperature (RT), shaking gently.
- 4) Rinse the membrane twice with PT-T20.
- 5) Incubate the membrane in anti-Caspase 8 [NB 500-208], diluted 1:500-1:1,000 in PT-T20 + 5% NFDM, for 60 minutes at RT.
- 6) Wash the membrane for 15 minutes, 3 times, in PT-T20 at RT.
- 7) Incubate the membrane in anti-mouse conjugated to HRP (secondary antibody), diluted in PT-T20 + 5% NFDM for 60 minutes at RT.
- 8) Wash the membrane for 15 minutes, 3 times, in PT-T20 at RT.
- 9) Develop membrane in chemiluminescent reagents, as instructed by kit-vendor.

*Positive control(s): Human kidney 293 cells





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB500-208

NB800-PC6	293 Whole Cell Lysate
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB500-208

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

