Product Datasheet

COPB2 Antibody (M3A5) NB600-102

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 10

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB600-102

Updated 5/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB600-102



NB600-102

COPB2 Antibody (M3A5)

COPB2 Antibody (M3A5)	
Product Information	
Unit Size	0.1 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	M3A5
Preservative	0.9% Sodium Azide
Isotype	IgG1
Purity	Unpurified
Buffer	Ascites
Product Description	
Host	Mouse
Gene ID	9276
Gene Symbol	COPB2
Species	Human, Rat, Avian, Bovine, Canine, Chicken, Hamster, Monkey, Primate, Rabbit
Reactivity Notes	kangaroo rat, goose
Specificity/Sensitivity	Recognizes an epitope shared by the beta-COP protein (110kD) found in most tissue culture cells, and a high MW doublet of brain microtubule-associated protein (MAP 2, 270-300 kD).
Immunogen	Microtubule-associated protein from goose brain.
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, Immunoaffinity Purification
Recommended Dilutions	Western Blot, Immunocytochemistry/ Immunofluorescence 1:20-1:50, Immunoprecipitation 1:10-1:500, Immunoaffinity Purification
Application Notes	AF: Use at an assay dependant dilution. ICC: Use at an assay dependant dilution. IF: CHO cells IP: Use at an assay dependant dilution. Stacked Golgi membranes from rat liver extract. Stains a reticular structure in the perinuclear area of non-neuronal cells (the periphery of the Golgi complex and a population of use idea assattant distribution and the sustained assattant dilution.



of vesicles scattered throughout the cytoplasm.

Publications

Schlatter A. On the Reality of Quantum Collapse and the Emergence of Space-Time Entropy 2019-03-25 [PMID: 33267037]

Schultz V, Barrie JA, Donald CL et al. Oligodendrocytes are susceptible to Zika virus infection in a mouse model of perinatal exposure: Implications for CNS complications Glia 2021-08-01 [PMID: 33942402]

BD Needham, M Funabashi, MD Adame, Z Wang, JC Boktor, J Haney, WL Wu, C Rabut, MS Ladinsky, SJ Hwang, Y Guo, Q Zhu, JA Griffiths, R Knight, PJ Bjorkman, MG Shapiro, DH Geschwind, DP Holschneid, MA Fischbach, SK Mazmanian A gut-derived metabolite alters brain activity and anxiety behaviour in mice Nature, 2022-02-14;602 (7898):647-653. 2022-02-14 [PMID: 35165440]

Ostrowski PP, Freeman SA, Fairn G, Grinstein S Dynamic Podosome-Like Structures in Nascent Phagosomes Are Coordinated by Phosphoinositides Dev. Cell 2019-06-06 [PMID: 31231039]

NakaYama H, Abe M, Morimoto C et al. Microglia permit climbing fiber elimination by promoting GABAergic inhibition in the developing cerebellum. Nat Commun. 2018-07-19 [PMID: 30026565]

McGugan, G et al. Characterization of a Rab11 -like GTPase, EhRab11, of Entamoeba histolytica. Molecular Biochemical Parasitology 1: 1-10. 2003-01-01 [PMID: 12850258]

Welter, B et al. Characterization of a Rab7-like GTPase, EhRab7: a marker for the earky stages of endocytosis in Entamoeba histolytica. Molecular Biochemical Parasitology 121:254-264. 2002-01-01 [PMID: 12034459]

Donaldson JG, Lippincott-Schwartz J, Klausner RD. Guanine nucleotides modulate the effects of brefeldin A in semipermeable cells: regulation of the association of a 110-kD peripheral membrane protein with the Golgi apparatus. J Cell Biol112(4):579-88. 1991-02-01 [PMID: 1993732] (ICC/IF, Bovine, Rat)

Donaldson JG, Lippincott-Schwartz J, Bloom GS et al. Dissociation of a 110-kD peripheral membrane protein from the Golgi apparatus is an early event in brefeldin A action. J Cell Biol111(6 Pt 1):2295-306. 1990-12-01 [PMID: 227706] (ICC/IF, Rat)

Cianciola NL, Carlin CR. Adenovirus RID-alpha activates an autonomous cholesterol regulatory mechanism that rescues defects linked to Niemann-Pick disease type C. J Cell Biol187(4):537-52. 2009-11-16 [PMID: 19948501]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB600-102

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)

H00009276-P01-10ug Recombinant Human COPB2 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-102

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

