

Product Datasheet

UCH-L1/PGP9.5 Antibody (31A3)

NB600-1160

Unit Size: 0.1 mg

Store at 4C.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 18

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB600-1160

Updated 4/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB600-1160




NB600-1160**UCH-L1/PGP9.5 Antibody (31A3)**

Product Information	
Unit Size	0.1 mg
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	31A3
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Product Description	
Description	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP2-33130) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	7345
Gene Symbol	UCHL1
Species	Human, Mouse, Rat, Porcine, Bovine, Canine, Zebrafish
Reactivity Notes	Zebrafish reactivity reported in scientific literature (PMID: 30377377).
Marker	pan-Neuronal Marker
Immunogen	UCH-L1/PGP9.5 protein from brain (Uniprot: P09936)
Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1-2 ug/ml, Simple Western 10 ug/ml, Flow Cytometry 1-2 ug/million cells, Immunohistochemistry 0.5-1.0ug/ml, Immunocytochemistry/Immunofluorescence 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Immunohistochemistry-Frozen 0.5-1.0ug/ml
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.




Images


Western Blot: UCH-L1/PGP9.5 Antibody (31A3) [NB600-1160] - Western Blot Analysis of human brain tissue lysate using UCH-L1/PGP9.5 Antibody (31A3)




Immunocytochemistry/Immunofluorescence: UCH-L1/PGP9.5 Antibody (31A3) [NB600-1160] - Immunofluorescence Analysis of T98G cells labeling Pgp9.5 with UCH-L1/PGP9.5 Antibody (31A3) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Nucspot (Red)




Immunohistochemistry-Paraffin: UCH-L1/PGP9.5 Antibody (31A3) [NB600-1160] - Formalin-fixed, paraffin-embedded human brain stained with UchL1 antibody (PGP9.5) (1:500), peroxidase-conjugate and DAB chromogen. Staining seen in cytoplasm, ER and membrane. Fixation in 95% ethanol/5% acetic acid for 2-3 hours prior to paraffin embedding is recommended. Specimens which have not been fixed in acetic acid/alcohol require pre-treatment with citrate buffer: for example, unmask of the target epitope by boiling the tissue sections in 10mM citrate buffer, pH 6.0, for 10 min followed by cooling at room temp for 20 min.




Flow Cytometry: UCH-L1/PGP9.5 Antibody (31A3) [NB600-1160] - Flow Cytometric Analysis of T98G cells using UCH-L1/PGP9.5 Antibody (31A3) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).



Western Blot: UCH-L1/PGP9.5 Antibody (31A3) [NB600-1160] - analysis of UchL1 in 1) human, 2) mouse and 3) rat brain lysate using UchL1 antibody at 1 ug/ml. goat anti-mouse Ig HRP secondary antibody and ECL substrate solution were used for this test.




Immunocytochemistry/Immunofluorescence: UCH-L1/PGP9.5 Antibody (31A3) [NB600-1160] - analysis of PGP9.5 in rat mesenteric artery using anti-PGP9.5 antibody. Image from verified customer review.



Simple Western: UCH-L1/PGP9.5 Antibody (31A3) [NB600-1160] - Simple Western lane view shows a specific band for PGP9.5 / UCHL-1 in 0.2 mg/ml of h. Cerebellum (left) and IMR-32 (right) lysate(s). This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Simple Western: UCH-L1/PGP9.5 Antibody (31A3) [NB600-1160] - Electropherogram images of the corresponding Simple Western lane. PGP9.5 / UCHL-1 antibody was used at 10 ug/ml dilution of h. Cerebellum and IMR-32 lysates(s) respectively.



Publications

Shi WG, Yao Y, Liang YJ et al. Activation of TGR5 in the injured nerve site according to a prevention protocol mitigates partial sciatic nerve ligation-induced neuropathic pain by alleviating neuroinflammation Pain 2024-10-25 [PMID: 39450924]

Kim CY, Lee EH, Kwak SH et al. UCHL1 Overexpression Is Related to the Aggressive Phenotype of Non-small Cell Lung Cancer Tuberculosis and Respiratory Diseases 2024-10-01 [PMID: 39362830]

Duan J, Lv A, Guo Z et al. CX3CR1 + /UCHL1 + microglial extracellular vesicles in blood: a potential biomarker for multiple sclerosis Journal of Neuroinflammation 2024-10-09 [PMID: 39385200]

Lucarini E, Micheli L, Pagnotta E et al. The Efficacy of Camelina sativa Defatted Seed Meal against Colitis-Induced Persistent Visceral Hypersensitivity: The Relevance of PPAR alpha Receptor Activation in Pain Relief Nutrients 2022-07-29 [PMID: 35956313] (IHC-P, Rat)

Details:

Dilution used 1:500

Manavis J, Blumbergs P, Jerrett I et al. Heterogeneous immunoreactivity of axonal spheroids in focal symmetrical encephalomalacia produced by Clostridium perfringens type D epsilon toxin in sheep Veterinary pathology 2021-12-06 [PMID: 34872408]

Sundaresan S, Meininger C A et al. Gastrin Induces Nuclear Export and Proteasome Degradation of Menin in Enteric Glial Cells. Gastroenterology 2017-01-12 [PMID: 28859856] (IF/IHC, ICC/IF, Mouse)

Pal P, Hales K, Hales DB Increased Transient Receptor Potential Melastatin 8 Expression in the development of bladder pain in patients with Interstitial cystitis/Bladder Pain Syndrome Oncotarget 2020-10-10 [PMID: 33088425] (ICC/IF, Human)

Moutal A, Ji Y, Bellampalli SS, Khanna R Differential expression of Cdk5-phosphorylated CRMP2 following a spared nerve injury Mol Brain 2020-06-22 [PMID: 32571373] (IF/IHC, Rat)

Arnaboldi F, Sommariva M, Opizzi E et al. Expression of Toll-like receptors 4 and 7 in murine peripheral nervous system development Ann. Anat. 2020-05-04 [PMID: 32380196] (ICC/IF, Mouse)

Alberio L, Locarno A, Saponaro A et al. A light-gated potassium channel for sustained neuronal inhibition. Nat. Methods. 2018-11-01 [PMID: 30377377] (IF/IHC, Zebrafish)

Novakovic P, Harding JC, Al-Dissi AN et al. Pathologic Evaluation of Type 2 Porcine Reproductive and Respiratory Syndrome Virus Infection at the Maternal-Fetal Interface of Late Gestation Pregnant Gilts PLoS ONE 2016-03-11 [PMID: 26963101] (IF/IHC, WB, Human)

Bottner M, Zorenkov D, Hellwig I et al. Expression pattern and localization of alpha-synuclein in the human enteric nervous system. Neurobiol Dis. 2012-12-01 [PMID: 22850485] (IHC-Fr)

Details:

IHC (frozen); Table I (colon carcinoma, dysplastic adenoma).

More publications at <http://www.novusbio.com/NB600-1160>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB600-1160

NBL1-17575	UCH-L1/PGP9.5 Overexpression Lysate
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-1160

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

