Product Datasheet

Vinculin Antibody (hVIN-1) NB600-1293

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 14

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB600-1293

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB600-1293



NB600-1293


Vinculin Antibody (hVIN-1)

Vinculin Antibody (hVIN-1)	
Product Information	
0.1 ml	
This product is unpurified. The exact concentration of antibody is not quantifiable.	
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Monoclonal	
hVIN-1	
0.9% Sodium Azide	
IgG1	
Unpurified	
Ascites	
116 kDa	
Mouse	
7414	
VCL	
Human, Mouse, Rat, Amphibian, Bovine, Canine, Chicken, Turkey	
Frog (100%). Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.	
Focal Adhesion Marker	
Specifically labels vinculin at cell-cell and cell-substrate contacts. Shows cross-reactivity with smooth muscle metavinculin.	
Purified human vinculin from uterus.	
Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Single Cell Western	
Western Blot 1:200 - 1:400, Simple Western, Immunohistochemistry 1:10 - 1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Frozen 1:10 - 1:500, Single Cell Western 1:30	
See <u>Simple Western Antibody Database</u> for Simple Western validation: tested in HeLa lysate; separated by size; antibody dilution of 1:5; matrix was 12-230 kDa. Single Cell Western reported by an internal validation on treated LNCap cells at a 1:30 dilution	




Images


Western Blot: Vinculin Antibody (hVIN-1) [NB600-1293] - Cell line lysates were separated on SDS-PAGE and probed with 1:200 Monoclonal Anti-Vinculin Clone: hVIN-1. The antibody was developed using Goat Anti-Mouse IgG-Peroxidase and a chemiluminescent substrate. Lanes: 1.HeLa 2.COS7 3.NIH-3T3 4.RAT2 5.CHO 6.MDBK 7.MDCK



Immunocytochemistry/Immunofluorescence: Vinculin Antibody (hVIN-1) [NB600-1293] - Immunofluorescence images of LM8 cells treated with 0 nM, 1 nM, or 10 nM eribulin and stained for vinculin (red) and nucleus (blue) (left). Dotted line shows the cell shape. Scale bar: 10 um. Quantitative analysis of the area of vinculin staining (right). Values are mean +/- SEM (less than or equal to 30 cells per group). **P < 0.01. Image collected and cropped by CiteAb from the following publication (//pubmed.ncbi.nlm.nih.gov/30719211/) licensed under a CC-BY license.



Immunohistochemistry: Vinculin Antibody (hVIN-1) [NB600-1293] - Enhanced Validation-By Independent Antibodies: Immunohistochemistry. Formalin-fixed, paraffin-embedded Rat Heart sections stained with 15 ug/mL Anti-Vinculin antibody produced in Rabbit (Cat. No. V4139) (A). The antibody was developed using Anti-Rabbit IgG (whole molecule)-FITC antibody produced in Goat (Cat. No. F9887), and 15 ug/mL Monoclonal Anti-Vinculin antibody produced in Mouse, Clone: hVIN1 (Cat. No. V9131) (B). The antibody was developed using Rabbit Anti-Mouse IgG-Cy3 conjugate antibody. Results: Two Anti-Vinculin antibodies, V4139 (A) and V9131 (B), target different regions of Vinculin show similar staining profiles between the two antibodies, demonstrating Independent Antibody Verification.



Publications

Hasegawa J, Nagata T, Ihara K et al. Heteroduplex oligonucleotide technology boosts oligonucleotide splice switching activity of morpholino oligomers in a Duchenne muscular dystrophy mouse model Nature Communications 2024-09-26 [PMID: 39327422]

X Zhou, S Wahane, MS Friedl, M Kluge, CC Friedel, K Avrampou, V Zachariou, L Guo, B Zhang, X He, RH Friedel, H Zou Microglia and macrophages promote corralling, wound compaction and recovery after spinal cord injury via Plexin-B2 Nat. Neurosci., 2020-03-01;23(3):337-350. 2020-03-01 [PMID: 32112058]

Li H, Huynh TN, Duong MT et al. ACAT1/SOAT1 Blockade Suppresses LPS-Mediated Neuroinflammation by Modulating the Fate of Toll-like Receptor 4 in Microglia International journal of molecular sciences 2023-03-15 [PMID: 36982689] (Western Blot, Mouse)

Ventura E, Xie C, Buraschi S et al. Complexity of progranulin mechanisms of action in mesothelioma Journal of experimental & clinical cancer research : CR 2022-12-05 [PMID: 36471440] (Immunocytochemistry/ Immunofluorescence, Human)

Li Y, Li C, Liu Q et al. Loss of Acta2 in cardiac fibroblasts does not prevent the myofibroblast differentiation or affect the cardiac repair after myocardial infarction Journal of molecular and cellular cardiology 2022-08-22 [PMID: 36007455] (IHC-Fr, Mouse)

Details:

IHC-Fr dilution 1:100

Wu L, Xu Y, Xi K et al. Regulation of macrophage subtype via injectable micro/nano-structured porous microsphere for reprogramming osteoimmune microenvironment Chemical Engineering Journal 2022-07-01 (ICC/IF, Rat)

Costanzo F, Martinez Diez M, Santamaria Nunez G et al. Promoters of ASCL1- and NEUROD1-dependent genes are specific targets of lurbinectedin in SCLC cells EMBO molecular medicine 2022-03-09 [PMID: 35263037] (WB, Human)

Uzureau S, Lecordier L, Uzureau P et Al. APOL1 C-Terminal Variants May Trigger Kidney Disease through Interference with APOL3 Control of Actomyosin Cell Rep 2020-03-17 [PMID: 32187552] (ICC/IF, Human)

Wu L, Gu Y, Liu L et al. Hierarchical micro/nanofibrous membranes of sustained releasing VEGF for periosteal regeneration Biomaterials 2019-10-18 [PMID: 31655445] (ICC/IF, Human)

Miao Q, Hill MC, Chen F et al. SOX11 and SOX4 drive the reactivation of an embryonic gene program during murine wound repair Nat Commun [PMID: 31492871] (ICC/IF, IF/IHC, Human)

Watanabe K, Yui Y, Sasagawa S et al. Low-dose eribulin reduces lung metastasis of osteosarcoma in vitro and in vivo Oncotarget 2019-01-04 [PMID: 30719211] (ICC/IF, Mouse)

Yadav N, Jaber FL, Sharma Y et al. Efficient reconstitution of hepatic microvasculature by endothelin receptor antagonism in liver sinusoidal endothelial cells. Hum. Gene Ther. 2018-09-28 [PMID: 30266073] (ICC/IF, Mouse)

More publications at http://www.novusbio.com/NB600-1293





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB600-1293

NBL1-17706 Vinculin Overexpression Lysate

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-1293

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

