

Product Datasheet

Cholesterol Oxidase Antibody - BSA Free NB600-625-100ug

Unit Size: 100 ug

Store lyophilized antibody at 4C. Aliquot reconstituted liquid and store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB600-625

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB600-625



NB600-625-100ug

Cholesterol Oxidase Antibody - BSA Free

Product Information	
Unit Size	100 ug
Concentration	LYOPH mg/ml
Storage	Store lyophilized antibody at 4C. Aliquot reconstituted liquid and store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.01% Sodium Azide
Reconstitution Instructions	Reconstitute with 100 ul deionized water (or equivalent)
Isotype	IgG
Purity	Multi-step
Buffer	Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Product Description	
Description	<p>This antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Cholesterol Oxidase [Microorganism]</p> <p>Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use.</p>
Host	Goat
Species	Bacteria
Reactivity Notes	Reactive with Cholesterol Oxidase from Streptomyces. Cross reactivity against Cholesterol Oxidase from other sources is unknown
Specificity/Sensitivity	Cholesterol Oxidase
Immunogen	Cholesterol Oxidase [Streptomyces] (Uniprot: P12676)
Product Application Details	
Applications	Western Blot, ELISA, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500-1:5000, ELISA 1:5000-1:20000, Immunoprecipitation 1:100
Application Notes	This product has been tested by western blot. This product is suitable to be assayed against 1.0 ug of Cholesterol Oxidase [Microorganism] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for this product.



Images


Western Blot: Cholesterol Oxidase Antibody [NB600-625] - Analysis using HRP conjugate of NB600-625. Detection of purified N-acylmanosamino-1-Dehydrogenase under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified target proteins contained 4% BME and were boiled for 5 minutes. Samples of 1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:3000 dilution of primary antibody. Detection shown was using Dylight 488 conjugated Donkey anti goat.



Western Blot: Cholesterol Oxidase Antibody [NB600-625] - Detection of purified cholesterol oxidase under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified cholesterol oxidase contained 4% BME and were boiled for 5 minutes. Samples of 1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with a 1:3000 dilution of primary antibody. Detection shown was using Dylight 488 conjugated Donkey anti goat.



Goat anti Cholesterol oxidase antibody (lot 6534) was used to detect purified cholesterol oxidase under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified cholesterol oxidase contained 4% BME and were boiled for 5 minutes. Samples of ~1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with a 1:3000 dilution of primary antibody (ON 4 C in). Detection shown was using Dylight 488 conjugated Donkey anti goat (lot 21094 1:10K in TBS/ 1 hr RT). Images were collected using the BioRad VersaDoc System.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB600-625-100ug

HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-625

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

