Product Datasheet

Goat anti-Mouse IgG1 Heavy Chain Secondary Antibody [FITC] NB7510

Unit Size: 1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB7510

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB7510



NB7510

Goat anti-Mouse IgG1 Heavy Chain Secondary Antibody [FITC]

Mouse IgG1

Immunohistochemistry-Paraffin

Soat anti-mouse igor meavy chain becondary Antibody [1110]	
Product Information	
Unit Size	1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C in the dark.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Conjugate	FITC
Purity	Immunogen affinity purified
Buffer	Phosphate Buffered Saline (PBS) containing 0.2%BSA
Product Description	
Host	Goat
Species	Mouse
Specificity/Sensitivity	By immunoelectrophoresis and ELISA this antibody reacts specifically with mouse IgG1. Cross reactivity to mouse IgA, IgM, IgG2a, IgG2b, IgG2c, IgG3 and IgE is undetectable. Some hybridoma clones may express aberrant

Publications

Product Application Details

Recommended Dilutions

Wang T, Gilkes DM, Takano N Et al. Hypoxia-inducible factors and RAB22A mediate formation of microvesicles that stimulate breast cancer invasion and metastasis Proc Natl Acad Sci U S A 2014-06-19 [PMID: 24938788] (ICC/IF)

1:50 - 1:500, Immunohistochemistry-Paraffin 1:50 - 1:500

immunoglobulin-related peptides that are improperly recognized by this antibody.

Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence

Immunocytochemistry/ Immunofluorescence, Immunohistochemistry,

Details:

Immunogen

Applications

Citation using the Cy3 version of this antibody.

Chaturvedi P, Gilkes DM, Takano N, Semenza GL. Hypoxia-inducible factor-dependent signaling between triplenegative breast cancer cells and mesenchymal stem cells promotes macrophage recruitment. Proc Natl Acad Sci U S A. [PMID: 24799675] (Chemotaxis)

Details:

Citation using the FITC form of this antibody.

Ochi T. Role of mitotic motors, dynein and kinesin, in the induction of abnormal centrosome integrity and multipolar spindles in cultured V79 cells exposed to dimethylarsinic acid. Mutat Res. [PMID: 11804606]

Details:

Citation using the FITC form of this antibody.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB7510

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

