Product Datasheet

Enolase 1 Overexpression Lysate NBL1-10267

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-10267

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-10267



NBL1-10267

Enolase 1 Overexpression Lysate


Unit Size0.1 mgConcentrationThe exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.StorageStore at -80.C. Avoid freeze-thaw cycles.BufferRIPA bufferTarget Molecular Weight47 kDaProduct DescriptionEnclase 1 Transient Overexpression LysatePrescriptionEnclase 1 Transient Overexpression LysatePission Host: HEK293TPlasmid: RC205494Accession#: NM_001428Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control to 100m DDT.Gene ID2023Gene SymbolENO1NotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volum DNT.NotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volum DNA transfection. Transfected cells were cultured for Alshis befor concentration was measured by BCA protein assay kit.Lysate TypeOverxpressionProduct Application DetailsWestern BlotApplicationsWestern BlotRecommended DilutionsWestern Blot	Product Information	
overexpression lysates. Please contact technical support for more information. Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 47 kDa Product Description Enclase 1 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC205494 Accession#: NM_001428 Protein Tag: C-MYC/DDK You will receive 1 vial of Lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml), The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Giycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 2023 Gene Symbol ENO1 Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for alkns before concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Overexpression	Unit Size	0.1 mg
Buffer RIPA buffer Target Molecular Weight 47 kDa Product Description Enclase 1 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC205494 Accession#: NM_001428 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 2023 Gene Symbol ENO1 Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection reagent specially designed and manufactured for large volume DNA transfection reagent specially designed to clarify the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Concentration	
Target Molecular Weight47 kDaProduct DescriptionEnclase 1 Transient Overexpression LysateDescriptionExpression Host: HEK293TPlasmid: RC205494Accession#: NM_001428Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample Buffer (250u). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID2023Gene SymbolHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3V04, and then centrifuged to calrify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailisWestern Blot	Storage	Store at -80C. Avoid freeze-thaw cycles.
Product Description Description Enclase 1 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC205494 Accession#: NM_001428 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 2023 Gene Symbol ENO1 Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3V04, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Buffer	RIPA buffer
DescriptionEnolase 1 Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC205494Accession#: NM_001428Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID2023Gene SymbolENO1SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were cultured for 48 hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3V04, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Target Molecular Weight	47 kDa
Enolase 1 Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC205494Accession#: NM_001428Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID2023Gene SymbolENO1SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were outlured for 48rrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl. 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3V04, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Product Description	
Accession#: NM_001428Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID2023Gene SymbolENO1SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfecte cells were ultured for 48ms before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Description	
Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID2023Gene SymbolENO1SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large polymer transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Plasmid: RC205494
You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID2023Gene SymbolENO1SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Accession#: NM_001428
(100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID2023Gene SymbolENO1SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Protein Tag: C-MYC/DDK
Gene SymbolENO1SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		(100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue,
SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Gene ID	2023
NotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot	Gene Symbol	ENO1
polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot	Species	Human
Product Application Details Applications Western Blot	Notes	polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein
Applications Western Blot	Lysate Type	Overexpression
	Product Application Details	
Recommended Dilutions Western Blot	Applications	Western Blot
	Recommended Dilutions	Western Blot



Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (<u>NBP1-71705</u>) present on the protein construct.
	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: Enolase 1 Overexpression Lysate (Adult Normal) [NBL1-10267] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for Enolase 1.



www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBL1-10267

NBP1-30262	Recombinant Human Enolase 1 Protein
7954-GM-010/CF	GM-CSF [Unconjugated]
H00002023-M01	Enolase 1 Antibody (8G8)
NB300-109	Tyrosine Hydroxylase Antibody

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-10267

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

