Product Datasheet

KERA Overexpression Lysate NBL1-12228

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-12228

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-12228



NBL1-12228

KERA Overexpression Lysate

Unit Size 0.1 mg Concentration The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information. Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 40.3 kDa Product Description Expression Host: HEK293T Description KERA Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC207509 Accession#: NM_007035 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control of usid of 2x8DS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 11081 Gene Symbol KERA Notes HeK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48ms before concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Product Information	
overexpression lysates. Please contact technical support for more information. Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 40.3 kDa Product Description Expression Hyster Description KERA Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC207509 Accession#: NM_007035 Protein Tag: C-MYC/DDK You will receive 1 vial of Lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml), The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 11081 Gene symbol KERA Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Unit Size	0.1 mg
Buffer RIPA buffer Target Molecular Weight 40.3 kDa Product Description Expression Lysate Description KERA Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC207509 Accession#: NM_007035 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 11081 Gene Symbol KERA Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for alrge volume DNA transfection reagent specially designed and manufactured for large volume DNA transfection reagent specially designed and manufactured for large volume DNA transfection reagent specially designed and manufactured for large volume DNA transfection reagent specially designed to clarify the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Concentration	
Target Molecular Weight 40.3 kDa Product Description Description KERA Transient Overexpression Lysate Expression Host: HEK293T Expression Host: HEK293T Plasmid: RC207509 Accession#: NM_007035 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 11081 Gene Symbol KERA Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1XProteinase inhibitor cocktail mix, 1mM PMSF and TMM Na3VO4, and then centifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Storage	Store at -80C. Avoid freeze-thaw cycles.
Product Description Description KERA Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC207509 Accession#: NM_007035 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 11081 Gene Symbol KERA Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3V04, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Buffer	RIPA buffer
Description KERA Transient Overexpression Lysate Expression Host: HEK293T Expression Host: HEK293T Plasmid: RC207509 Accession#: NM_007035 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 11081 Gene Symbol KERA Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Mestern Blot	Target Molecular Weight	40.3 kDa
KERA Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC207509Accession#: NM_007035Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID11081Gene SymbolKERASpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were outured for 48rs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3V04, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Product Description	
Plasmid: RC207509Accession#: NM_007035Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID11081Gene SymbolKERASpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for darks before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Description	KERA Transient Overexpression Lysate
Accession#: NM_007035Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID11081Gene SymbolKERASpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot		Expression Host: HEK293T
Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID11081Gene SymbolKERASpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Plasmid: RC207509
You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID11081Gene SymbolKERASpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Accession#: NM_007035
(100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID11081Gene SymbolKERASpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Protein Tag: C-MYC/DDK
Gene SymbolKERASpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot		(100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue,
SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot	Gene ID	11081
NotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Gene Symbol	KERA
polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot	Species	Human
Product Application Details Applications Western Blot	Notes	polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein
Applications Western Blot	Lysate Type	Overexpression
	Product Application Details	
Recommended Dilutions Western Blot	Applications	Western Blot
	Recommended Dilutions	Western Blot



Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (<u>NBP1-71705</u>) present on the protein construct.
	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: KERA Overexpression Lysate (Adult Normal) [NBL1-12228] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for KERA.

158-	
106 - 79 -	
48 -	
35- 23-	
10	

www.novusbio.com



technical@novusbio.com



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBL1-12228

NBP1-84425PEP	KERA Recombinant Protein Antigen
233-FB-025	FGF basic/FGF2/bFGF [Unconjugated]
10838-KE-050	KERA [Unconjugated]
NB300-223	Vimentin Antibody

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-12228

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

