Product Datasheet

MMP-3 Overexpression Lysate NBL1-13159

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-13159

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-13159



NBL1-13159

MMP-3 Overexpression Lysate

Product Information				
Unit Size	0.1 mg			
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.			
Storage	Store at -80C. Avoid freeze-thaw cycles.			
Buffer	RIPA buffer			
Target Molecular Weight	52.2 kDa			
Product Description				
Description	MMP-3 Transient Overexpression Lysate			
	Expression Host: HEK293T			
	Plasmid: RC210235			
	Accession#: NM_002422			
	Protein Tag: C-MYC/DDK			
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.			
Gene ID	4314			
Gene Symbol	MMP3			
Species	Human			
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.			
Lysate Type	Overexpression			
Product Application Details				
Applications	Western Blot			
Recommended Dilutions	Western Blot			



Ea for de	LAG) epitope tag (<u>NBP1-71705</u>) present on the protein construct.
an	ich vial of cell lysate contains 100ug of total protein which should be sufficient 20-50 reactions. Depending on over-expression level, antibody affinity and tection system, some lysates can go as low as 0.1 ug per load. We commend starting with 5ug of cell lysate. Add an equal amount of cell lysate d 2X SDS Sample buffer and boil the SDS samples for 10 minutes before ading.

Images

Western Blot: MMP3 Overexpression Lysate (Adult Normal) [NBL1-13159] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for MMP3.

158-		
106-		
79 —		
48 —	-	
35-		
35- 23-		





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBL1-13159

NB100-91878PEP	MMP-3 Antibody Blocking Peptide
210-TA-005	TNF-alpha [Unconjugated]
DMP300	MMP-3 [HRP]
M6000B-1	IL-6 [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-13159

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

