Product Datasheet

PRELP Overexpression Lysate NBL1-14748

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-14748

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-14748



NBL1-14748

PRELP Overexpression Lysate

Product Information	
Init Size 0.	.1 mg
	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.
Storage S	Store at -80C. Avoid freeze-thaw cycles.
Buffer R	RIPA buffer
arget Molecular Weight 4	1.6 kDa
Product Description	
Pescription	PRELP Transient Overexpression Lysate
E	Expression Host: HEK293T
Ρ	Plasmid: RC207609
А	Accession#: NM_201348
Р	Protein Tag: C-MYC/DDK
(1 cc of	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control 100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate ontains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 00mM DTT.
Sene ID 55	549
Gene Symbol P	PRELP
pecies H	luman
ם ענ כנ 1! P	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid objective transfection reagent specially designed and manufactured for large olume DNA transfection. Transfected cells were cultured for 48hrs before ollection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 50mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.
ysate Type O	Overexpression
Product Application Details	
Applications V	Vestern Blot
Recommended Dilutions	Vestern Blot



Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.	Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (<u>NBP1-71705</u>) present on the protein construct.
		for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before

Images

Western Blot: PRELP Overexpression Lysate (Adult Normal) [NBL1-14748] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for PRELP.

158-	N. 1 2.2
106-	
79 —	
48 —	-
35- 23-	and the second
23 -	
10	Contraction of the second





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBL1-14748

AF808	Osteopontin/OPN Antibody [Unconjugated]
AF6447	PRELP Antibody [Unconjugated]
NBP1-91258	Fibronectin Antibody - BSA Free
NB100-74350	Aggrecan Neoepitope Antibody - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-14748

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

