Product Datasheet

PSMB5 Overexpression Lysate NBL1-14880

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-14880

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-14880



NBL1-14880

PSMB5 Overexpression Lysate

Concentration The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information. Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 28.3 kDa Product Description Psints Description PSMB5 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC209326 Accession#: NM_002797 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250u). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 5693 Gene symbol PSMB5 Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25MM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, intm EDTA, 1*Proteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3V04, and then centrifuged to clarify the lysate. Protein concontration was measu	Product Information			
overexpression lysates. Please contact technical support for more information.StorageStore at -80C. Avoid freeze-thaw cycles.BufferRIPA bufferTarget Molecular Weight28.3 kDaProduct DescriptionPSMB5 Transient Overexpression LysateDescriptionPSMB5 Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC209326Accession#: NM_002797Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5693Gene SymbolPSMB5SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (250mL Tris-HCl pH7.6, collection. The cells were lysed in modified RIPA buffer (250mL Tris-HCl pH7.6, collection. The cells were lysed in modified RIPA buffer (250mL Tris-HCl pH7.6, collection. The cells were lysed in modified RIPA buffer (250mL Tris-HCl pH7.6, collection. The cells were lysed in modified RIPA buffer PCL pH7.6, collection. The cells were lysed in modified RIPA buffer PCL pH7.6, collection. The cells were lysed in modified RIPA buffer PCL pH7.6, collection. The cells were lysed in modified RIPA buffer PCL pH7.6, collection. The cells were lysed in modified RIPA buffer PCL pH7.6, collection. The cells were lysed in modified RIPA buffer PCL pH7.6, collection. The cells were lysed in modified RIPA	Unit Size	0.1 mg		
Buffer RIPA buffer Target Molecular Weight 28.3 kDa Product Description Psime State Description PSMB5 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC209326 Accession#: NM_002797 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 5693 Gene Symbol PSMB5 Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection reagent specially designed and manufactured for large volume DNA transfection reagent specially designed to clarity the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Concentration			
Target Molecular Weight 28.3 kDa Product Description PSMB5 Transient Overexpression Lysate Expression Host: HEK293T Expression Host: HEK293T Plasmid: RC209326 Accession#: NM_002797 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250u). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 5693 Gene Symbol PSMB5 Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl PH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1XProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VQ4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Storage	Store at -80C. Avoid freeze-thaw cycles.		
Product Description Description PSMB5 Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC209326 Accession#: NM_002797 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250u). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 5693 Gene Symbol PSMB5 Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were tor large to rolem A 1mM Na3VQ4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Buffer	RIPA buffer		
Description PSMB5 Transient Overexpression Lysate Expression Host: HEK293T Expression Host: HEK293T Plasmid: RC209326 Accession#: NM_002797 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT. Gene ID 5693 Gene Symbol PSMB5 Species Human Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for large volume DNA transfection reagent specially designed and manufacture of Protein accentration was measured by BCA protein assay kit. Lysate Type Overexpression Product Application Details Western Blot	Target Molecular Weight	28.3 kDa		
PSMB5 Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC209326Accession#: NM_002797Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSD5 sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5693Gene SymbolPSMB5SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were outlured for large volume DNA transfection. Transfected cells were cultured for 48ns before collection. The cells were lysed in modified RIPA buffer (25MM Tris-HCl pH7.6, 150mM NaCl. 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3V04, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Product Description			
Plasmid: RC209326Plasmid: RC209326Accession#: NM_002797Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5693Gene SymbolPSMB5SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were eultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot	Description	PSMB5 Transient Overexpression Lysate		
Accession#: NM_002797Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5693Gene SymbolPSMB5SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Expression Host: HEK293T		
Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5693Gene SymbolPSMB5SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Plasmid: RC209326		
You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5693Gene SymbolPSMB5SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot		Accession#: NM_002797		
(100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5693Gene SymbolPSMB5SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application DetailsWestern Blot		Protein Tag: C-MYC/DDK		
Gene SymbolPSMB5SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot		(100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue,		
SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot	Gene ID	5693		
NotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot	Gene Symbol	PSMB5		
polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details ApplicationsWestern Blot	Species	Human		
Product Application Details Applications Western Blot	Notes	polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein		
Applications Western Blot	Lysate Type	Overexpression		
	Product Application Details			
Recommended Dilutions Western Blot	Applications	Western Blot		
	Recommended Dilutions	Western Blot		



Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (<u>NBP1-71705</u>) present on the protein construct.
	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: PSMB5 Overexpression Lysate (Adult Normal) [NBL1-14880] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for PSMB5.

158-	28	
106 - 79 -		
79 -		
48 —		
35-		
35- 23-		-
		100





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBL1-14880

NBP1-50934-0.05mg	Recombinant Human PSMB5 His Protein
AF114	CD45 Antibody [Unconjugated]
NBP2-13820	PSMB5 Antibody
NB600-717	MBP Antibody (12)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-14880

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

