Product Datasheet

PGD2 Synthase/PTGDS Overexpression Lysate NBL1-14930

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-14930

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-14930



PGD2 Synthase/PTGDS Ove	rexpression Lysate
Product Information	
Unit Size	0.1 mg
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.
Storage	Store at -80C. Avoid freeze-thaw cycles.
Buffer	RIPA buffer
Target Molecular Weight	20.8 kDa
Product Description	
Description	PGD2 Synthase/PTGDS Transient Overexpression Lysate
	Expression Host: HEK293T
	Plasmid: RC216795
	Accession#: NM_000954
	Protein Tag: C-MYC/DDK
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.
Gene ID	5730
Gene Symbol	PTGDS
Species	Human
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid

ore FI pH7.6, hix, 1mM otein
large

Lysate Type	Overexpression
Lysale Type	Overexpression

Product Application Details		
Applications	Western Blot	
Recommended Dilutions	Western Blot	




Application Notes

This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (NBP1-71705) present on the protein construct.

Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: Prostaglandin D Synthase/Lipocalin Overexpression Lysate (Adult Normal) [NBL1-14930] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for Prostaglandin D Synthase/Lipocalin.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBL1-14930

NB110-59909PEP PGD2 Synthase/PTGDS Antibody Blocking Peptide

7954-GM-010/CF GM-CSF [Unconjugated]

NBP1-79280 PGD2 Synthase/PTGDS Antibody

NB100-689 COX-2 Antibody

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-14930

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

