## **Product Datasheet**

### SNX16 Overexpression Lysate NBL1-16308

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-16308

Updated 11/9/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-16308



#### NBL1-16308

SNX16 Overexpression Lysate

| roduct Information         |   |
|----------------------------|---|
| Init Size 0                | ).1 mg  |
|                            | The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.  |
| storage S                  | Store at -80C. Avoid freeze-thaw cycles.  |
| Ruffer R                   | RIPA buffer   |
| arget Molecular Weight 3   | 39 kDa  |
| Product Description        |   |
|                            | SNX16 Transient Overexpression Lysate   |
| E                          | Expression Host: HEK293T  |
| Ρ                          | Plasmid: RC216565   |
| A                          | Accession#: NM_022133   |
| Ρ                          | Protein Tag: C-MYC/DDK  |
| (1<br>Ci                   | You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control<br>100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate<br>contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists<br>of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue,<br>00mM DTT.   |
| iene ID 6                  | 64089   |
| Gene Symbol S              | SNX16   |
| pecies H                   | luman   |
| P<br>Vi<br>Ci<br>1<br>P    | HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid<br>polymer transfection reagent specially designed and manufactured for large<br>volume DNA transfection. Transfected cells were cultured for 48hrs before<br>collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6,<br>50mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM<br>PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein<br>concentration was measured by BCA protein assay kit. |
| ysate Type C               | Dverexpression  |
| roduct Application Details |   |
| pplications V              | Vestern Blot  |
|                            |   |



| Application Notes | This product is intended for use as a positive control in Western Blot.<br>Overexpression of the target protein was confirmed using an antibody to DDK<br>(FLAG) epitope tag ( <u>NBP1-71705</u> ) present on the protein construct.   |
|-------------------|--|
|                   | Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading. |

#### Images

| Western Blot: SNX16 Overexpression Lysate (Adult Normal) [NBL1-<br>16308] Left-Empty vector transfected control cell lysate (HEK293 cell<br>lysate); Right -Over-expression Lysate for SNX16. | 158 -<br>106 - |
|---|----------------|
|   | 79             |
|   | 48-            |
|   | 35-<br>23-     |
| Western Blot: SNX16 Overexpression Lysate (Adult Normal) [NBL1-<br>16308] - Western Blot experiments. Left-Control; Right -Over-expres  | 158-           |
| Lysate for SNX16  | 106 —<br>79 —  |
|   | 48 -           |
|   | 35 -<br>23 -   |
|   | 25             |





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBL1-16308

| H00064089-P01-10ug | Recombinant Human SNX16 GST (N-Term) Protein |
|--------------------|--|
| 236-EG-200         | EGF [Unconjugated]                           |
| NB100-1211         | SNX16 Antibody                               |
| AF467              | EphB2 Antibody [Unconjugated]                |
|                    |  |

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-16308

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

